arXiv:1108.0460v4 [math.FA] 25 Jul 2012

α-Modulation Spaces (I) Scaling, Embedding and Algebraic Properties

Jinsheng Han
Department of Mathematics, Shanghai Jiao Tong University, Shanghai 200240, P. R. China.
Baoxiang Wang
LMAM, School of Mathematics, Peking University and BICMR, Beijing 100871, P. R. China.
(pkuhan@sjtu.edu.cn, wbx@pku.edu.cn)

Abstract. First, we consider some fundamental properties including dual spaces, complex interpolations of α-modulation spaces $M^{s,\alpha}_{p,q}$ with $0 < p, q \le \infty$. Next, necessary and sufficient conditions for the scaling property and the inclusions between α_1-modulation and α_2-modulation spaces are obtained. Finally, we give some criteria for α-modulation spaces constituting multiplication algebra. As a by-product, we show that there exists an α-modulation space which is not an interpolation space between modulation and Besov spaces. In a subsequent paper, we will give some applications of α-modulation spaces to nonlinear dispersive wave equations.

Key words. α-modulation space, Dual space, Multiplication algebra, Scaling property, Embedding.

AMS subject classifications. 42 B35, 42 B37, 35 A23

1 Introduction and definition

Frequency localization technique plays an important role in the modern theory of function spaces. There are two kinds of basic partitions to the Euclidean space $\mathbb{R}^n$, one is the dyadic decomposition $\mathbb{R}^n = \{\xi : |\xi| < 1\} \bigcup (\bigcup_{j=1}^{\infty} \{\xi : |\xi| \in [2^{j-1}, 2^j)\})$, another is the uniform decomposition $\mathbb{R}^n = \bigcup_{k\in\mathbb{Z}^n} (k+[-1/2, 1/2)^n)$. According to these two kinds of decompositions in frequency spaces, one can naturally introduce the dyadic decomposition operators $\triangle_j$ ($j \in \mathbb{Z}_+$) whose symbol φ_j is localized in $\{\xi : |\xi| \sim 2^j\}$, and the uniform decomposition operator $\Box_k$ ($k \in \mathbb{Z}^n$) whose symbol σ_k is supported in $k + [-1, 1]^n$. The difference between φ_j and σ_k is that the diameters of supp φ_j and supp σ_k are $O(2^j)$ and $O(1)$, respectively. All tempered distributions acted on these decomposition operators with finite $\ell^q(L^p)$ (quasi)-norms constitute Besov space $B^s_{p,q}$ and modulation space $M^s_{p,q}$, respectively.

The α-modulation spaces $M^{s,\alpha}_{p,q}$, introduced by Gröbner [11], are proposed to be intermediate function spaces to connect modulation space and Besov space with respect to parameters $\alpha \in [0, 1]$, which are formulated by some new kind of α-decomposition operators $\Box^{\alpha}_k$ ($k \in \mathbb{Z}^n$). We denote by η^{α}_k the symbol of $\Box^{\alpha}_k$, whose essential characteristic is that the diameter of its support set has power growth as $\langle k\rangle^{\alpha/(1-\alpha)}$.

Modulation spaces are special α-modulation spaces in the case $\alpha = 0$, and Besov space can be regarded as the limit case of α-modulation space when $\alpha \nearrow 1$. Modulation spaces were first introduced by Feichtinger [8] in the study of time-frequency analysis to consider the decay property of a function in both physical and frequency spaces. His original idea is to use the short-time Fourier transform of a tempered distribution equipping with a mixed $L^q(L^p)$-norm to generate modulation spaces $M^s_{p,q}$. Gröchenig's book [12] systematically discussed the theory of time-frequency analysis and modulation spaces. In Gröbner's doctoral thesis, he used the α-covering to the frequency space $\mathbb{R}^n$ and a corresponding bounded admissible partition of unity of order p (p-BAPU) to define α-modulation spaces. Some recent works have been devoted to the study of α-modulation spaces (see [1, 2, 7, 10, 14, 13] and references therein). Borup and Nielsen [1] and Fornasier [10] constructed Banach frames for α-modulation spaces in the multivariate setting, Kobayashi, Sugimoto and Tomita [14, 13] discussed the boundedness for a class of pseudo-differential operators with symbols in α-modulation spaces. Dahlke, Fornasier, Rauhut, Steidl and Teschke [7] established the relationship between the generalized coorbit theory and α-modulation spaces. The aim of the present paper is to describe some standard properties including the dual spaces, embeddings, scaling and algebraic structure of α-modulation spaces.

Before stating the notion of α-modulation spaces, we introduce some notations frequently used in this paper. $A \lesssim B$ stands for $A \le CB$, and $A \sim B$ denote $A \lesssim B$ and $B \lesssim A$, where C is a positive constant which can be different at different places. Let $\mathscr{S}(\mathbb{R}^n)$ be the Schwartz space and $\mathscr{S}'(\mathbb{R}^n)$ be its strongly topological dual space. Suppose $f \in \mathscr{S}'(\mathbb{R}^n)$ and $\lambda > 0$, we write $f_\lambda(\cdot) = f(\lambda\cdot)$. Let X be a (quasi-)Banach space, we denote by X^* the dual space of X. For any $p \in [1,\infty]$, p^* will stand for the dual number of p, i.e., $1/p + 1/p^* = 1$. We denote by $L^p = L^p(\mathbb{R}^n)$ the Lebesgue space for which the norm is written by $\|\cdot\|_p$, and by ℓ^p the sequence Lebesgue space. We will write $\langle x\rangle = (1+|x|^2)^{1/2}$. For any multi-index $\delta = (\delta_1, \delta_2, \cdots, \delta_n)$, we denote $D^\delta = \partial_1^{\delta_1}\partial_2^{\delta_2}\cdots\partial_n^{\delta_n}$. It is convenient to divide $\mathbb{R}^n$ into n parts $\mathbb{R}^n_j, j = 0, 1, 2, \cdots, n$:

$$\mathbb{R}^n_j = \{x \in \mathbb{R}^n : |x_i| \leqslant |x_j|,\ i = 1, \cdots, j-1, j+1, \cdots, n\}.$$

We write $J = (I-\Delta)^{s/2}$ and define the Sobolev space

$$H^s(\mathbb{R}^n) = \left\{f \in \mathscr{S}'(\mathbb{R}^n) : \|f\|_{H^s} = \|J^s f\|_2 < \infty\right\}$$

and

$$\ell^q_{s,\alpha}(\mathbb{Z}^n; L^p) = \left\{\{g_k\}_{k\in\mathbb{Z}^n} : g_k \in \mathscr{S}'(\mathbb{R}^n), \left\|\langle k\rangle^{\frac{s}{1-\alpha}}\|g_k\|_p\right\|_{\ell^q} < \infty\right\}.$$

Without additional note, we will always assume that

$$s \in \mathbb{R}, \quad 0 < p, q \leqslant \infty, \quad 0 \leqslant \alpha < 1.$$

Let us start with the third partition of unity on frequency space for $\alpha \in [0,1)$ (see [1]). We suppose $c < 1$ and $C > 1$ are two positive constants, which relate to the space dimension n, and a Schwartz function sequence $\{\eta^\alpha_k\}_{k\in\mathbb{Z}^n}$ satisfies

$$|\eta^\alpha_k(\xi)| \gtrsim 1, \quad \text{if } \left|\xi - \langle k\rangle^{\frac{\alpha}{1-\alpha}}k\right| < c\langle k\rangle^{\frac{\alpha}{1-\alpha}}; \tag{1.1a}$$

$$\mathrm{supp}\,\eta^\alpha_k \subset \left\{\xi : \left|\xi - \langle k\rangle^{\frac{\alpha}{1-\alpha}}k\right| < C\langle k\rangle^{\frac{\alpha}{1-\alpha}}\right\}; \tag{1.1b}$$

$$\sum\nolimits_{k\in\mathbb{Z}^n} \eta^\alpha_k(\xi) \equiv 1, \quad \forall \xi \in \mathbb{R}^n; \tag{1.1c}$$

$$\langle k\rangle^{\frac{\alpha|\delta|}{1-\alpha}}\left|D^\delta \eta^\alpha_k(\xi)\right| \lesssim 1, \quad \forall \xi \in \mathbb{R}^n. \tag{1.1d}$$

We denote

$$\Upsilon = \left\{ \{\eta_k^\alpha\}_{k\in\mathbb{Z}^n} : \{\eta_k^\alpha\}_{k\in\mathbb{Z}^n} \text{ satisfies (1.1)} \right\} \tag{1.2}$$

Corresponding to every sequence $\{\eta_k^\alpha\}_{k\in\mathbb{Z}^n} \in \Upsilon$, one can construct an operator sequence denoted by $\{\Box_k^\alpha\}_{k\in\mathbb{Z}^n}$, and

$$\Box_k^\alpha = \mathscr{F}^{-1}\eta_k^\alpha\mathscr{F}. \tag{1.3}$$

Υ is nonempty. Indeed, let ρ be a smooth radial bump function supported in $B(0,2)$, satisfying $\rho(\xi)=1$ as $|\xi|<1$, and $\rho(\xi)=0$ as $|\xi|\geqslant 2$. For any $k\in\mathbb{Z}^n$, we set

$$\rho_k^\alpha(\xi) = \rho\left(\frac{\xi - \langle k\rangle^{\frac{\alpha}{1-\alpha}}k}{C\langle k\rangle^{\frac{\alpha}{1-\alpha}}}\right) \tag{1.4}$$

and denote

$$\eta_k^\alpha(\xi) = \rho_k^\alpha(\xi)\left(\sum_{l\in\mathbb{Z}^n}\rho_l^\alpha(\xi)\right)^{-1}.$$

It is easy to verify that $\{\eta_k^\alpha\}_{k\in\mathbb{Z}^n}$ satisfies (1.1). This type of decomposition on frequency space is a generalization of the uniform decomposition and the dyadic decomposition. When $0\leqslant\alpha<1$, on the basis of this decomposition, we define the α-modulation space by

$$M_{p,q}^{s,\alpha}(\mathbb{R}^n) = \left\{ f\in\mathscr{S}'(\mathbb{R}^n) : \|f\|_{M_{p,q}^{s,\alpha}} = \|\{\Box_k^\alpha f\}_{k\in\mathbb{Z}^n}\|_{\ell_{s,\alpha}^q(\mathbb{Z}^n;L^p)} < \infty \right\}. \tag{1.5}$$

Denote $\varphi(\xi)=\rho(\xi)-\rho(2\xi)$, we may assume $\varphi(\xi)=1$ if $5/8\leq|\xi|\leq 3/2$. We introduce the function sequence $\{\varphi_k\}_{k=0}^\infty$:

$$\begin{cases} \varphi_j(\xi) = \varphi(2^{-j}\xi), \quad j\in\mathbb{N}, \\ \varphi_0(\xi) = 1 - \sum_{j=1}^\infty \varphi_j(\xi). \end{cases} \tag{1.6}$$

Define

$$\triangle_j = \mathscr{F}^{-1}\varphi_j\mathscr{F}, \quad j\in\mathbb{N}\cup\{0\}, \tag{1.7}$$

$\{\triangle_j\}_{j=0}^\infty$ is said to be the Littlewood–Paley (or dyadic) decomposition operators. Denote

$$B_{p,q}^s(\mathbb{R}^n) = \left\{ f\in\mathscr{S}'(\mathbb{R}^n) : \|f\|_{B_{p,q}^s} = \left\|\{2^{sj}\triangle_j f\}_{j\in\mathbb{N}\cup\{0\}}\right\|_{\ell^q(L^p)} < \infty \right\}. \tag{1.8}$$

Strictly speaking, (1.5) cannot cover the case $\alpha=1$, however, we will denote $M_{p,q}^{s,1}=B_{p,q}^s$ for convenience.

The paper is organized as follows. In Section 2, we show some basic properties on α-modulation spaces, their dual and complex interpolation spaces are presented there. In Section 3, we discuss the scaling property. In Section 4, the inclusions between α-modulation spaces for different indices α (including Besov spaces) are obtained. In Section 5, we study the regularity conditions so that α-modulation spaces form multiplication algebra. Finally, we show the necessity for the conditions of scalings, embeddings and algebra structures by constructing several counterexamples.

2 Some basic properties

In the sequel, we give some basic properties of $M^{s,\alpha}_{p,q}$. We need the following

Proposition 2.1 (**[18], Convolution in L^p with $p < 1$**). *Let $0 < p \leq 1$. $L^p_{B(x_0,R)} = \{f \in L^p(\mathbb{R}^n) : \operatorname{supp} f \subset B(x_0, R)\}$, $B(x_0, R) = \{x : |x - x_0| \leq R\}$. Suppose that $f, g \in L^p_{B(x_0,R)}$, then there exists a constant $C > 0$ which is independent of x_0 and $R > 0$ such that*

$$\|f * g\|_p \leq CR^{n(1/p-1)}\|f\|_p\|g\|_p.$$

Proposition 2.2. (**[18], Generalized Bernstein inequality**) *Let $\Omega \subset \mathbb{R}^n$ be a compact set, $0 < r \leq \infty$. Let us denote $\sigma_r = n(1/(r \wedge 1) - 1/2)$ and assume that $s > \sigma_r$. Then there exists a constant $C > 0$ such that*

$$\|\mathscr{F}^{-1}\varphi\mathscr{F}f\|_r \leq C\|\varphi\|_{H^s}\|f\|_r \tag{2.1}$$

holds for all $f \in L^r_\Omega := \{f \in L^p : \operatorname{supp}\widehat{f} \subset \Omega\}$ and $\varphi \in H^s$. Moreover, if $r \geq 1$, then (2.1) *holds for all $f \in L^r$.*

Proposition 2.3 (**Equivalent norm**). *Let $\{\eta^\alpha_k\}_{k\in\mathbb{Z}^n}, \{\widetilde{\eta}^\alpha_k\}_{k\in\mathbb{Z}^n} \in \Upsilon$, then they genarate equivalent quasi-norms on $M^{s,\alpha}_{p,q}$.*

Proof. See [1]. ☐

Proposition 2.4 (**Embedding**). *Let $0 < p_1 \leqslant p_2 \leq \infty$, $0 < q_1, q_2 \leq \infty$. We have*
(i) if $q_1 \leqslant q_2$ and $s_1 \geqslant s_2 + n\alpha\big(\frac{1}{p_1} - \frac{1}{p_2}\big)$, then

$$M^{s_1,\alpha}_{p_1,q_1} \subset M^{s_2,\alpha}_{p_2,q_2}; \tag{2.2}$$

(ii) if $q_1 > q_2$ and $s_1 > s_2 + n\alpha\big(\frac{1}{p_1} - \frac{1}{p_2}\big) + n(1-\alpha)\big(\frac{1}{q_2} - \frac{1}{q_1}\big)$, then

$$M^{s_1,\alpha}_{p_1,q_1} \subset M^{s_2,\alpha}_{p_2,q_2}. \tag{2.3}$$

Proof. From Bernstein's inequality it follows that

$$\|\square^\alpha_k f\|_{p_2} \lesssim \langle k\rangle^{\frac{n\alpha}{1-\alpha}\left(\frac{1}{p_1}-\frac{1}{p_2}\right)}\|\square^\alpha_k f\|_{p_1}. \tag{2.4}$$

Then (i) follows from $\ell^{q_1} \subset \ell^{q_2}$ and (2.4). For (ii), we use Hölder's inequality to obtain

$$\|f\|_{M^{s_2,\alpha}_{p_2,q_2}} \lesssim \|\{\square^\alpha_k f\}_{k\in\mathbb{Z}^n}\|_{\ell^{q_1}_{s_1,\alpha}}\|\{1\}_{k\in\mathbb{Z}^n}\|_{\ell^{q_1q_2/(q_1-q_2)}_{s_2-s_1+n\alpha(1/p_1-1/p_2),\alpha}}.$$

For the second term in the right-hand side, we easily see that it is finite by changing the summation to an integration. ☐

Proposition 2.5 (**Completeness**). *(i) $M^{s,\alpha}_{p,q}$ is a quasi-Banach space, and is a Banach space if $1 \leqslant p \leqslant \infty$ and $1 \leqslant q \leqslant \infty$.*
(ii) We have

$$\mathscr{S}(\mathbb{R}^n) \subset M^{s,\alpha}_{p,q}(\mathbb{R}^n) \subset \mathscr{S}'(\mathbb{R}^n). \tag{2.5}$$

Moreover, if $0 < p, q < \infty$, then $\mathscr{S}(\mathbb{R}^n)$ is dense in $M^{s,\alpha}_{p,q}$.

Proof. See [3]. ☐

Proposition 2.6 (Isomorphism). *For any $\sigma \in \mathbb{R}$, the mapping $J^\sigma : M^{s,\alpha}_{p,q} \to M^{s-\sigma,\alpha}_{p,q}$ is isomorphic.*

Proof. See [18] $\square$

Proposition 2.7. *$M^{s,\alpha}_{2,2}(\mathbb{R}^n) = H^s(\mathbb{R}^n)$ with equivalent norms.*

Proof. Plancherel's identity implies the result, see [11]. $\square$

2.1 Duality

It is known that the dual space of Besov space $B^s_{p,q}$ is $B^{-s+n(1/(p\wedge 1)-1)}_{(p\vee 1)^*,(q\vee 1)^*}$ (see [18]) and the dual space of modulation space $M^s_{p,q}$ is $M^{-s}_{(p\vee 1)^*,(q\vee 1)^*}$ (see [19]). In this section we study the dual spaces of α-modulation spaces.

Proposition 2.8. *Suppose $1 \leqslant p, q < \infty$. Then we have*

$$\left(\ell^q_{s,\alpha}(\mathbb{Z}^n; L^p)\right)^* = \ell^{q^*}_{-s,\alpha}(\mathbb{Z}^n; L^{p^*}).$$

More precisely, $f \in \left(\ell^q_{s,\alpha}(\mathbb{Z}^n; L^p)\right)^*$ is equivalent to that there exists a sequence $\{f_k\}_{k\in\mathbb{Z}^n} \in \ell^{q^*}_{-s,\alpha}(\mathbb{Z}^n; L^{p^*})$ such that for any $g = \{g_k\}_{k\in\mathbb{Z}^n} \in \ell^q_{s,\alpha}(\mathbb{Z}^n; L^p)$, we have

$$\langle f, g\rangle = \sum_{k\in\mathbb{Z}^n} \int_{\mathbb{R}^n} f_k(x)\overline{g_k(x)}dx,$$

with $\|f\|_{(\ell^q_{s,\alpha}(\mathbb{Z}^n;L^p))^*} = \|\{f_k\}\|_{\ell^{q^*}_{-s,\alpha}(\mathbb{Z}^n;L^{p^*})}$.

It is a direct consequence of Proposition 3.3 in [19].

Lemma 2.1. *Let $\{g_k\}_{k\in\mathbb{Z}^n} \in \ell^q_{s,\alpha}(\mathbb{Z}^n; L^p)$, and Γ be a subset of $\mathbb{Z}^n$, then we have*

$$\Big\|\sum_{k\in\Gamma} \square^\alpha_k g_k\Big\|_{M^{s,\alpha}_{p,q}} \lesssim \|\{\square^\alpha_k g_k\}_{k\in\Gamma}\|_{\ell^q_{s,\alpha}(\mathbb{Z}^n;L^p)}. \tag{2.6}$$

Proof. We introduce

$$\Lambda(k) = \{l \in \mathbb{Z}^n : \square^\alpha_k \square^\alpha_l f \neq 0\}. \tag{2.7}$$

We denote the constant in (1.1b) relating to $\{\eta^\alpha_l\}_{l\in\mathbb{Z}^n}$ by C, thus for every $l \in \Lambda(k)$, there holds

$$\langle k\rangle^{\frac{\alpha}{1-\alpha}}(k_j - C) < \langle l\rangle^{\frac{\alpha}{1-\alpha}}(l_j + C), \tag{2.8a}$$

$$\langle k\rangle^{\frac{\alpha}{1-\alpha}}(k_j + C) > \langle l\rangle^{\frac{\alpha}{1-\alpha}}(l_j - C) \tag{2.8b}$$

with $j = 1, \cdots, n$. For the above l and k, we conclude that

$$\langle k\rangle \sim \langle l\rangle. \tag{2.9}$$

If $|k| \lesssim 1$ (or $|l| \lesssim 1$), it is easy to see that (2.9) follows from (2.8). Thus, it suffices to show (2.9) in the case $|k| \gg 1$ (or $|l| \gg 1$). When $k \in \mathbb{R}^n_j$ with $k_j > 0$, from (2.8a); whereas when $k \in \mathbb{R}^n_j$ but with $k_j < 0$, from (2.8b)$\times(-1)$, we see $\langle k\rangle^{\frac{1}{1-\alpha}} \lesssim \langle l\rangle^{\frac{1}{1-\alpha}}$, and symmetrically, we have $\langle l\rangle^{\frac{1}{1-\alpha}} \lesssim \langle k\rangle^{\frac{1}{1-\alpha}}$. Therefore, we get (2.9). Suppose both l and $\widetilde{l}$ are in $\Lambda(k)$. substituting l with $\widetilde{l}$ (2.8a) and (2.8b) also hold. It follows that

$$\left|\langle l\rangle^{\frac{\alpha}{1-\alpha}} l_j - \langle \widetilde{l}\rangle^{\frac{\alpha}{1-\alpha}} \widetilde{l}_j\right| \lesssim \langle k\rangle^{\frac{\alpha}{1-\alpha}} + \langle l\rangle^{\frac{\alpha}{1-\alpha}} + \langle \widetilde{l}\rangle^{\frac{\alpha}{1-\alpha}}.$$

Then Taylor's theorem, combined with (2.9), gives $|l_j - \widetilde{l}_j| \lesssim 1$. It follows that

$$\#\Lambda(k) \sim 1. \tag{2.10}$$

One has that the right hand side of (2.6) is

$$\begin{aligned}\left(\sum_{l\in\mathbb{Z}^n} \langle l\rangle^{\frac{sq}{1-\alpha}} \left\| \sum_{k\in\Lambda(l)\cap\Gamma} \Box_l^\alpha \Box_k^\alpha g_k \right\|_p^q\right)^{\frac{1}{q}} &\lesssim \left(\sum_{l\in\mathbb{Z}^n} \langle l\rangle^{\frac{sq}{1-\alpha}} \sum_{k\in\Lambda(l)\cap\Gamma} \|\Box_l^\alpha \Box_k^\alpha g_k\|_p^q\right)^{\frac{1}{q}} \\ &\lesssim \left(\sum_{k\in\Gamma} \|\Box_k^\alpha g_k\|_p^q \sum_{l\in\Lambda(k)} \langle l\rangle^{\frac{sq}{1-\alpha}}\right)^{\frac{1}{q}} \\ &\lesssim \left(\sum_{k\in\Gamma} \langle k\rangle^{\frac{sq}{1-\alpha}} \|\Box_k^\alpha g_k\|_p^q\right)^{\frac{1}{q}}.\end{aligned} \tag{2.11}$$

We remark that in the second inequality of (2.11), by Young's inequality, or by Proposition 2.1 we see that

$$\|\Box_k^\alpha \Box_l^\alpha f\|_p \lesssim \|\Box_k^\alpha f\|_p,$$

which enable us to remove $\Box_l^\alpha$; and in the third inequality of (2.11), we have applied (2.10) to remove the summation on $l \in \Lambda(k)$. □

Theorem 2.1. *Suppose* $0 < p, q < \infty$*, then we have*

$$\left(M_{p,q}^{s,\alpha}\right)^* = M_{(1\vee p)^*,(1\vee q)^*}^{-s+n\alpha\left(\frac{1}{1\wedge p}-1\right)}. \tag{2.12}$$

Proof. The proof is separated into four cases.

Case 1: $1 \leqslant p, q < \infty$. First, we show that $\left(M_{p,q}^{s,\alpha}\right)^* \subset M_{p^*,q^*}^{-s,\alpha}$. Noticing that

$$M_{p,q}^{s,\alpha} \ni f \to \{\Box_k^\alpha f\} \in \ell_{s,\alpha}^q(\mathbb{Z}^n; L^p)$$

is an isometric mapping from $M_{p,q}^{s,\alpha}$ onto a subspace $X = \{\{\Box_k^\alpha f\} : f \in M_{p,q}^{s,\alpha}\}$ of $\ell_{s,\alpha}^q(\mathbb{Z}^n; L^p)$, so, any continuous functional g on $M_{p,q}^{s,\alpha}$ can be regarded as a bounded linear functional on X, which can be extended onto $\ell_{s,\alpha}^q(\mathbb{Z}^n; L^p)$ (the extension is written as $\tilde{g}$) and the norm of g is preserved. By Proposition 2.8, there exists $\{g_k\} \in \ell_{-s,\alpha}^{q^*}(\mathbb{Z}^n; L^{p^*})$ such that

$$\langle \tilde{g}, \{f_k\}\rangle = \sum_{k\in\mathbb{Z}^n} \int \overline{g_k(x)} f_k(x) dx \tag{2.13}$$

holds for all $\{f_k\} \in \ell_{s,\alpha}^q(\mathbb{Z}^n; L^p)$. Moreover, $\|g\|_{(M_{p,q}^{s,\alpha})^*} = \|\{g_k\}\|_{\ell_{-s,\alpha}^{q^*}(\mathbb{Z}^n; L^{p^*})}$. Since $M_{p,q}^{s,\alpha}$ is isometric to X, we see that

$$\langle g, \varphi\rangle = \langle \tilde{g}, \{\Box_k^\alpha \varphi\}\rangle = \int \sum_{k\in\mathbb{Z}^n} \overline{\Box_k^\alpha g_k(x)} \varphi(x) dx, \quad \varphi \in M_{p,q}^{s,\alpha}, \tag{2.14}$$

Hence, $g = \sum_{k\in\mathbb{Z}^n} \Box_k^\alpha g_k(x)$. In view of Lemma 2.1 and Young's inequality,

$$\|g\|_{M^{-s,\alpha}_{p^*,q^*}} \lesssim \|\{g_k\}_{k\in\mathbb{Z}^n}\|_{\ell^{q^*}_{-s,\alpha}(\mathbb{Z}^n;L^{p^*})} = \|g\|_{(M^{s,\alpha}_{p,q})^*},$$

which implies $\left(M^{s,\alpha}_{p,q}\right)^* \subset M^{-s,\alpha}_{p^*,q^*}$. Next, we prove the reverse inclusion. For any $f \in M^{-s,\alpha}_{p^*,q^*} \subset \mathscr{S}'$, we show that $f \in \left(M^{s,\alpha}_{p,q}\right)^*$. Let $\varphi \in \mathscr{S}$. We have

$$\begin{aligned}|\langle f,\varphi\rangle| &\leqslant \|\{\Box_k^\alpha f\}_{k\in\mathbb{Z}^n}\|_{\ell^{q^*}_{-s,\alpha}(\mathbb{Z}^n;L^{p^*})} \left\| \left\{ \sum_{l\in\Lambda(k)} \Box_l^\alpha \varphi \right\}_{k\in\mathbb{Z}^n} \right\|_{\ell^q_{s,\alpha}(\mathbb{Z}^n;L^p)} \\ &\lesssim \|f\|_{M^{-s,\alpha}_{p^*,q^*}} \|\varphi\|_{M^{s,\alpha}_{p,q}}.\end{aligned}$$

The principle of duality implies $M^{-s,\alpha}_{p^*,q^*} \subset \left(M^{s,\alpha}_{p,q}\right)^*$.

In the following, we discuss the left three cases. From (2.2) in Proposition 2.4, we know

$$M^{s,\alpha}_{p,q} \subset M^{s-n\alpha\left(\frac{1}{p\wedge 1}-1\right),\alpha}_{1\vee p,1\vee q}.$$

This combined with the principle of duality gives

$$\left(M^{s,\alpha}_{p,q}\right)^* \supset \left(M^{s-n\alpha\left(\frac{1}{p\wedge 1}-1\right),\alpha}_{1\vee p,1\vee q}\right)^* = M^{-s+n\alpha\left(\frac{1}{p\wedge 1}-1\right),\alpha}_{(1\vee p)^*,(1\vee q)^*}.$$

Hence, only the reverse inclusion needs to be proven.

Case 2. $1 \leqslant p < \infty, 0 < q < 1$. For any $f \in \left(M^{s,\alpha}_{p,q}\right)^*$, take any $k \in \mathbb{Z}^n$ and any $\varphi \in \mathscr{S}(\mathbb{R}^n)$, we have

$$\begin{aligned}|\langle \Box_k^\alpha f,\varphi\rangle| &= |\langle f, \Box_k^\alpha \varphi\rangle| \\ &\leqslant \|f\|_{\left(M^{s,\alpha}_{p,q}\right)^*} \|\Box_k^\alpha \varphi\|_{M^{s,\alpha}_{p,q}} \\ &\lesssim \langle k\rangle^{\frac{s}{1-\alpha}} \|f\|_{\left(M^{s,\alpha}_{p,q}\right)^*} \|\varphi\|_p.\end{aligned} \tag{2.15}$$

This implies $\left(M^{s,\alpha}_{p,q}\right)^* \subset M^{-s,\alpha}_{p^*,\infty}$.

Case 3. $0 < p, q < 1$. For any $f \in \left(M^{s,\alpha}_{p,q}\right)^*$, take any $k \in \mathbb{Z}^n$, we have

$$\begin{aligned}|\Box_k^\alpha f(x)| &= |\langle f, \overline{\mathscr{F}^{-1}\eta_k^\alpha(x-\cdot)}\rangle| \\ &= |\langle f, \mathscr{F}^{-1}\eta_k^\alpha(\cdot - x)\rangle| \\ &\lesssim \|f\|_{\left(M^{s,\alpha}_{p,q}\right)^*} \|\mathscr{F}^{-1}\eta_k^\alpha(\cdot - x)\|_{M^{s,\alpha}_{p,q}} \\ &\lesssim \langle k\rangle^{\frac{s}{1-\alpha}-\frac{n\alpha}{1-\alpha}\left(\frac{1}{p}-1\right)} \|f\|_{\left(M^{s,\alpha}_{p,q}\right)^*}.\end{aligned} \tag{2.16}$$

This implies $\left(M^{s,\alpha}_{p,q}\right)^* \subset M^{-s+n\alpha\left(\frac{1}{p}-1\right),\alpha}_{\infty,\infty}$.

Case 4. $0 < p < 1, 1 \leqslant q < \infty$. For any $f \in \left(M^{s,\alpha}_{p,q}\right)^*$ and every $k \in \mathbb{Z}^n$, there exists some $x_k \in \mathbb{R}^n$ satisfying

$$\|\Box_k^\alpha f\|_\infty \sim |\mathscr{F}^{-1}\eta_k^\alpha \mathscr{F} f(x_k)|.$$

Let $\{a_k\}$ be an arbitrary $\ell^q_{s-n\alpha(1/p-1)}$ sequence, and we construct another sequence namely $\{\widetilde{a}_k\}$, such that $|\widetilde{a}_k| = |a_k|$, and the argument of $\widetilde{a}_k$ is the opposite number of the principal argument of $\mathscr{F}^{-1}\eta_k^\alpha \mathscr{F} f(x_k)$. From (1.1b),(1.1d), we get $\|\mathscr{F}^{-1}\eta_k^\alpha(\cdot - x_k)\|_p \lesssim \langle k\rangle^{-\frac{n\alpha}{1-\alpha}\left(\frac{1}{p}-1\right)}$. Therefore, we have

$$\begin{aligned}\sum_{k\in\mathbb{Z}^n}|a_k|\|\Box_k^\alpha f\|_\infty &\sim \left\langle f, \sum_k \widetilde{a}_k \mathscr{F}^{-1}\eta_k^\alpha(\cdot - x_k)\right\rangle \\ &\leqslant \|f\|_{\left(M_{p,q}^{s,\alpha}\right)^*}\left\|\sum_k \widetilde{a}_k \mathscr{F}^{-1}\eta_k^\alpha(\cdot - x_k)\right\|_{M_{p,q}^{s,\alpha}} \\ &\lesssim \|f\|_{\left(M_{p,q}^{s,\alpha}\right)^*}\|\{a_k\}_{k\in\mathbb{Z}^n}\|_{\ell^q_{s-n\alpha(\frac{1}{p}-1)}}.\end{aligned}$$

This implies $\left(M_{p,q}^{s,\alpha}\right)^* \subset M_{\infty,q^*}^{-s+n\alpha\left(\frac{1}{p}-1\right),\alpha}$. □

2.2 Complex interpolation

The complex interpolation for Besov spaces has a beautiful theory; cf. [18]. We can imitate the counterpart for the Besov space to construct the complex interpolation for α-modulation spaces. It will be repeatedly used in the following argument. Since there is little essential modification in the statement, we only provide the outline of the proof.

We start with some abstract theory about complex interpolation on quasi-Banach spaces. Let $S = \{z : 0 < \mathrm{Re} z < 1\}$ be a strip in the complex plane. Its closure $\{z : 0 \leqslant \mathrm{Re} z \leqslant 1\}$ is denoted by $\overline{S}$. We say that $f(z)$ is an $\mathscr{S}'(\mathbb{R}^n)$-analytic function in S if the following properties are satisfied:
(i) for every fixed $z \in \overline{S}$, $f(z) \in \mathscr{S}'(\mathbb{R}^n)$;
(ii) for any $\varphi \in \mathscr{S}(\mathbb{R}^n)$ with compact support, $\mathscr{F}^{-1}\varphi\mathscr{F} f(x,z)$ is a uniformly continuous and bounded function in $\mathbb{R}^n \times \overline{S}$;
(iii) for any $\varphi \in \mathscr{S}(\mathbb{R}^n)$ with compact support, $\mathscr{F}^{-1}\varphi\mathscr{F} f(x,z)$ is an analytic function in S for every fixed $x \in \mathbb{R}^n$.
We denote the set of all $\mathscr{S}'(\mathbb{R}^n)$-analytic functions in S by $\mathbf{A}(\mathscr{S}'(\mathbb{R}^n))$. The idea we used here is due to Calderón [4], Calderón and Torchinsky [5, 6] and Triebel [18].

Definition 2.1. *Let A_0 and A_1 be quasi-Banach spaces, and $0 < \theta < 1$. We define*

$$\begin{aligned}\mathbf{F}(A_0, A_1) =& \Big\{\varphi(z) \in \mathbf{A}(\mathscr{S}'(\mathbb{R}^n)) : \varphi(\ell + it) \in A_\ell, \ell = 0, 1, \forall t \in \mathbb{R}, \\ & \|\varphi(z)\|_{\mathbf{F}(A_0,A_1)} \overset{\triangle}{=} \max_{\ell=0,1}\sup_{t\in\mathbb{R}}\|\varphi(\ell+it)\|_{A_\ell}\Big\};\end{aligned} \tag{2.17}$$

and

$$\begin{aligned}(A_0, A_1)_\theta =& \Big\{f \in \mathscr{S}'(\mathbb{R}^n) : \exists\varphi(z) \in \mathbf{F}(A_0, A_1) \text{ such that } f = \varphi(\theta), \\ & \|f\|_{(A_0,A_1)_\theta} \overset{\triangle}{=} \inf_\varphi \|\varphi(z)\|_{\mathbf{F}(A_0,A_1)}\Big\},\end{aligned} \tag{2.18}$$

where the infimum is taken over all $\varphi(z) \in \mathbf{F}(A_0, A_1)$ such that $\varphi(\theta) = f$.

The following two propositions are essentially known in [18] and the references therein.

Proposition 2.9. *Suppose all notations have the same meaning as in Definition 2.1, then we have*

$$\big((A_0, A_1)_\theta, \|\cdot\|_{(A_0,A_1)_\theta}\big)$$

is a quasi-Banach space.

Proposition 2.10. *Suppose all notations have the same meaning as in Definition 2.1, then we have*

$$\|f\|_{(A_0,A_1)_\theta} = \inf_{\varphi}\left(\sup_{t\in\mathbb{R}}\|\varphi(it)\|_{A_0}^{1-\theta}\sup_{t\in\mathbb{R}}\|\varphi(1+it)\|_{A_1}^{\theta}\right), \tag{2.19}$$

where the infimum is taken over all $\varphi(z) \in \mathbf{F}(A_0, A_1)$ such that $\varphi(\theta) = f$.

We point out the interpolation functor referred in (2.18) is an exact interpolation functor of exponent θ. For our purpose, we will use the following multi-linear case.

Proposition 2.11. *Let T be a continuous multi-linear operator from $A_0^{(1)} \times A_0^{(2)} \times \cdots \times A_0^{(m)}$ to B_0 and from $A_1^{(1)} \times A_1^{(2)} \times \cdots \times A_1^{(m)}$ to B_1, satisfying*

$$\begin{aligned}
&\big\|T\big(f^{(1)}, f^{(2)}, \cdots, f^{(m)}\big)\big\|_{B_0} \leqslant C_0\prod_{j=1}^{m}\|f^{(j)}\|_{A_0^{(j)}};\\
&\big\|T\big(f^{(1)}, f^{(2)}, \cdots, f^{(m)}\big)\big\|_{B_1} \leqslant C_1\prod_{j=1}^{m}\|f^{(1)}\|_{A_1^{(1)}},
\end{aligned} \qquad f^{(j)} \in A_0^{(j)} \cap A_1^{(j)}.$$

Then T is continuous from $\big(A_0^{(1)}, A_1^{(1)}\big)_\theta \times \big(A_0^{(2)}, A_1^{(2)}\big)_\theta \times \cdots \times \big(A_0^{(m)}, A_1^{(m)}\big)_\theta$ to $(B_0, B_1)_\theta$ with norm at most $C_0^{1-\theta}C_1^{\theta}$, provided $0 \leqslant \theta \leqslant 1$.

Proof. From Proposition 2.10, we know there exist m sequences $\{\varphi_k^{(j)}(z)\}_{k\in\mathbb{N}}, j = 1, \cdots, m$ satisfying

$$\lim_{k\to\infty}\sup_t\|\varphi_k^{(j)}(\mathrm{i}t)\|_{A_0}^{1-\theta}\sup_t\|\varphi_k^{(j)}(1+\mathrm{i}t)\|_{A_1}^{\theta} = \|f^{(j)}\|_{(A_0,A_1)_\theta}. \tag{2.20}$$

We put $\psi_k^{(j)}(z) = C_0^{\frac{z-1}{m}}C_1^{-\frac{z}{m}}T(\varphi_k^{(j)}(z))$. It is easy to see that $\psi_k^{(j)}(z) \in F(B_0, B_1)$ with $\psi_k^{(j)}(\theta) = C_0^{\theta-1}C_1^{-\theta}Tf^{(j)}$, and

$$\|\psi_k^{(j)}(\ell+\mathrm{i}t)\|_{B_\ell} \leqslant C_\ell^{-\frac{1}{m}}\|T\varphi_k^{(j)}(\ell+\mathrm{i}t)\|_{B_\ell} \leqslant \|\varphi_k^{(j)}(\ell+\mathrm{i}t)\|_{A_\ell}, \quad \ell = 0, 1.$$

Thus, combining Proposition 2.10, we have

$$\begin{aligned}
\big\|T\big(f^{(1)}, \cdots, f^{(m)}\big)\big\|_{(B_0,B_1)_\theta} &= C_0^{1-\theta}C_1^{\theta}\prod_{j=1}^{m}\|\psi_k^{(j)}(\theta)\|_{(B_0,B_1)_\theta}\\
&\leqslant C_0^{1-\theta}C_1^{\theta}\prod_{j=1}^{m}\left(\sup_t\|\psi_k^{(j)}(\mathrm{i}t)\|_{B_0}^{1-\theta}\sup_t\|\psi_k^{(j)}(1+\mathrm{i}t)\|_{B_1}^{\theta}\right) \qquad (2.21)\\
&\leqslant C_0^{1-\theta}C_1^{\theta}\prod_{j=1}^{m}\left(\sup_t\|\varphi_k^{(j)}(\mathrm{i}t)\|_{A_0}^{1-\theta}\sup_t\|\varphi_k^{(j)}(1+\mathrm{i}t)\|_{B_1}^{\theta}\right)
\end{aligned}$$

The conclusion follows from (2.21),(2.20). $\square$

Theorem 2.2. *Suppose* $0 < \theta < 1$ *and*

$$s = (1-\theta)s_0 + \theta s_1, \quad \frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \quad \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}, \tag{2.22}$$

then we have

$$\left(M^{s_0,\alpha}_{p_0,q_0}, M^{s_1,\alpha}_{p_1,q_1}\right)_\theta = M^{s,\alpha}_{p,q}. \tag{2.23}$$

Sketch of Proof. For $z \in \overline{S}$, we write

$$s(z) = (1-z)s_0 + zs_1, \quad \frac{1}{p(z)} = \frac{1-z}{p_0} + \frac{z}{p_1}, \quad \frac{1}{q(z)} = \frac{1-z}{q_0} + \frac{z}{q_1}.$$

For any $f \in M^{s,\alpha}_{p,q}$, we set

$$\varphi(x,z) = \sum_{k \in \mathbb{Z}^n} \langle k \rangle^{\frac{1}{1-\alpha}\left[\frac{sq}{q(z)} - s(z)\right]} \|\Box^\alpha_k f\|_p^{\frac{q}{q(z)} - \frac{p}{p(z)}} (\Box^\alpha_k f)^{\frac{p}{p(z)}}(x).$$

Obviously, $\varphi(z) \in \mathbf{A}(\mathscr{S}'(\mathbb{R}^n))$ and $\varphi(\theta) = f$. Direct calculation shows

$$\|\varphi(\ell + \mathrm{i}t)\|_{M^{s_\ell,\alpha}_{p_\ell,q_\ell}} \lesssim \|f\|_{M^{s,\alpha}_{p,q}}, \quad \ell = 0, 1.$$

This proves that $M^{s,\alpha}_{p,q} \subset \left(M^{s_0,\alpha}_{p_0,q_0}, M^{s_1,\alpha}_{p_1,q_1}\right)_\theta$.

Conversely, for any $f \in \left(M^{s_0,\alpha}_{p_0,q_0}, M^{s_1,\alpha}_{p_1,q_1}\right)_\theta$, if $\varphi \in \mathbf{A}(\mathscr{S}'(\mathbb{R}^n))$ such that $\varphi(\theta) = f$, for some $\theta \in (0,1)$, we can find two positive functions $\mu_0(\theta,t)$ and $\mu_1(\theta,t)$ in $(0,1) \times \mathbb{R}$ satisfying

$$\|\Box^\alpha_k f\|_p \leqslant \left(\frac{1}{1-\theta}\int_{\mathbb{R}} \|\Box^\alpha_k \varphi(\mathrm{i}t)\|^r_{p_0} \mu_0(\theta,t)dt\right)^{\frac{1-\theta}{r}} \left(\frac{1}{\theta}\int_{\mathbb{R}} \|\Box^\alpha_k \varphi(1+\mathrm{i}t)\|^r_{p_1} \mu_1(\theta,t)dt\right)^{\frac{\theta}{r}},$$

with $\frac{1}{1-\theta}\int_{\mathbb{R}} \mu_0(\theta,t)dt = \frac{1}{\theta}\int_{\mathbb{R}} \mu_1(\theta,t)dt = 1$. Taking the $\ell^q_{s,\alpha}$ norm of both sides leads to

$$\begin{aligned} \|\{\Box^\alpha_k f\}_{k \in \mathbb{Z}^n}\|_{\ell^q_{s,\alpha}} \leqslant{} & \left\| \frac{1}{1-\theta}\int_{\mathbb{R}} \langle k \rangle^{\frac{s_0 r}{1-\alpha}} \|\Box^\alpha_k \varphi(\mathrm{i}t)\|^r_{p_0} \mu_0(\theta,t)dt \right\|^{\frac{1-\theta}{r}}_{\ell^{\frac{q_0}{r}}} \\ & \times \left\| \frac{1}{\theta}\int_{\mathbb{R}} \langle k \rangle^{\frac{s_1 r}{1-\alpha}} \|\Box^\alpha_k \varphi(1+\mathrm{i}t)\|^r_{p_1} \mu_1(\theta,t)dt \right\|^{\frac{\theta}{r}}_{\ell^{\frac{q_1}{r}}}. \end{aligned} \tag{2.24}$$

Then, Minkowski's inequality implies that

$$\|f\|_{M^{s,\alpha}_{p,q}} \lesssim \sup_t \|\varphi(\mathrm{i}t)\|_{M^{s_0,\alpha}_{p_0,q_0}} \sup_t \|\varphi(1+\mathrm{i}t)\|_{M^{s_1,\alpha}_{p_1,q_1}}.$$

This proves $\left(M^{s_0,\alpha}_{p_0,q_0}, M^{s_1,\alpha}_{p_1,q_1}\right)_\theta \subset M^{s,\alpha}_{p,q}$. □

The following is a natural consequence of Proposition 2.11 and Theorem 2.2, and is frequently used later on.

Corrolary 2.1. *Suppose* T *is a continuous multi-linear mapping from* $M^{s^{(1)}_0,\alpha}_{p^{(1)}_0,q^{(1)}_0} \times \cdots \times M^{s^{(m)}_0,\alpha}_{p^{(m)}_0,q^{(m)}_0}$ *to* $M^{s_0,\alpha}_{p_0,q_0}$ *with norm* M_0*, and is also continuous, multi-linear from* $M^{s^{(1)}_1,\alpha}_{p^{(1)}_1,q^{(1)}_1} \times \cdots \times M^{s^{(m)}_1,\alpha}_{p^{(m)}_1,q^{(m)}_1}$ *to* $M^{s_1,\alpha}_{p_1,q_1}$ *with norm* M_1*. Then* T *is continuous and multi-linear from* $M^{s^{(1)},\alpha}_{p^{(1)},q^{(1)}} \times \cdots \times M^{s^{(m)},\alpha}_{p^{(m)},q^{(m)}}$ *to* $M^{s,\alpha}_{p,q}$ *with norm at most* $M_0^{1-\theta}M_1^\theta$*, provided* $0 \leqslant \theta \leqslant 1$*, and*

$$s^{(j)} = (1-\theta)s^{(j)}_0 + \theta s^{(j)}_1, \quad \frac{1}{p^{(j)}} = \frac{1-\theta}{p^{(j)}_0} + \frac{\theta}{p^{(j)}_1}, \quad \frac{1}{q^{(j)}} = \frac{1-\theta}{q^{(j)}_0} + \frac{\theta}{q^{(j)}_1}, \quad j = 1 \cdots m.$$

3 Scaling property

For Besov space, it is well known that

$$\|f_\lambda\|_{B^s_{p,q}} \lesssim \lambda^{-\frac{n}{p}}(1 \vee \lambda^s)\|f\|_{B^s_{p,q}}. \tag{3.1}$$

For modulation spaces with $s=0$ and $1 \le p,q \le \infty$, the sharp dilation property was obtained in [15] and they showed

$$\|f_\lambda\|_{M^0_{p,q}} \lesssim \lambda^{-\frac{n}{p}} \lambda^{0 \vee n\left(\frac{1}{q}-\frac{1}{p}\right) \vee n\left(\frac{1}{p}+\frac{1}{q}-1\right)} \|f\|_{M^0_{p,q}}, \quad \lambda > 1; \tag{3.2}$$

$$\|f_\lambda\|_{M^0_{p,q}} \lesssim \lambda^{-\frac{n}{p}} \lambda^{-\left[0 \vee n\left(\frac{1}{p}-\frac{1}{q}\right) \vee n\left(1-\frac{1}{p}-\frac{1}{q}\right)\right]} \|f\|_{M^0_{p,q}}, \quad \lambda < 1. \tag{3.3}$$

In this section, we study the scaling property of α-modulation spaces. For $0 < p,q \leqslant \infty$ and $(\alpha_1,\alpha_2) \in [0,1] \times [0,1]$, we denote

$$R(p,q;\alpha_1,\alpha_2) = 0 \vee \left[n(\alpha_1-\alpha_2)\left(\tfrac{1}{q}-\tfrac{1}{p}\right)\right] \vee \left[n(\alpha_1-\alpha_2)\left(\tfrac{1}{p}+\tfrac{1}{q}-1\right)\right], \tag{3.4}$$

which will be frequently used in this and the next sections. Then, we divide $\mathbb{R}^2_+$ into 3 sub-domains in two ways (see Fig. 1). One way is, $\mathbb{R}^2_+ = \mathrm{S}_1 \cup \mathrm{S}_2 \cup \mathrm{S}_3$ with

$$\begin{aligned}
&\mathrm{S}_1 = \left\{\left(\tfrac{1}{p},\tfrac{1}{q}\right) \in \mathbb{R}^2_+ : \tfrac{1}{q} \geqslant \tfrac{1}{p}, \tfrac{1}{p} \leqslant \tfrac{1}{2}\right\};\\
&\mathrm{S}_2 = \left\{\left(\tfrac{1}{p},\tfrac{1}{q}\right) \in \mathbb{R}^2_+ : \tfrac{1}{p}+\tfrac{1}{q} \geqslant 1, \tfrac{1}{p} > \tfrac{1}{2}\right\};\\
&\mathrm{S}_3 = \mathbb{R}^2_+ \backslash \{\mathrm{S}_1 \cup \mathrm{S}_2\},
\end{aligned}$$

Another way is, $\mathbb{R}^2_+ = \mathrm{T}_1 \cup \mathrm{T}_2 \cup \mathrm{T}_3$ with

$$\begin{aligned}
&\mathrm{T}_1 = \left\{\left(\tfrac{1}{p},\tfrac{1}{q}\right) \in \mathbb{R}^2_+ : \tfrac{1}{p} \geqslant \tfrac{1}{q}, \tfrac{1}{p} > \tfrac{1}{2}\right\};\\
&\mathrm{T}_2 = \left\{\left(\tfrac{1}{p},\tfrac{1}{q}\right) \in \mathbb{R}^2_+ : \tfrac{1}{p}+\tfrac{1}{q} \leqslant 1, \tfrac{1}{p} \leqslant \tfrac{1}{2}\right\};\\
&\mathrm{T}_3 = \mathbb{R}^2_+ \backslash \{\mathrm{T}_1 \cup \mathrm{T}_2\}.
\end{aligned}$$

If $\alpha_1 \geqslant \alpha_2$, then

$$R(p,q;\alpha_1,\alpha_2) = \begin{cases} n(\alpha_1-\alpha_2)\left(\frac{1}{q}-\frac{1}{p}\right), & \left(\frac{1}{p},\frac{1}{q}\right) \in \mathrm{S}_1;\\ n(\alpha_1-\alpha_2)\left(\frac{1}{p}+\frac{1}{q}-1\right), & \left(\frac{1}{p},\frac{1}{q}\right) \in \mathrm{S}_2;\\ 0, & \left(\frac{1}{p},\frac{1}{q}\right) \in \mathrm{S}_3. \end{cases} \tag{3.5}$$

If $\alpha_1 < \alpha_2$, then

$$R(p,q;\alpha_1,\alpha_2) = \begin{cases} 0, & \left(\frac{1}{p},\frac{1}{q}\right) \in \mathrm{T}_3;\\ n(\alpha_1-\alpha_2)\left(\frac{1}{p}+\frac{1}{q}-1\right), & \left(\frac{1}{p},\frac{1}{q}\right) \in \mathrm{T}_2;\\ n(\alpha_1-\alpha_2)\left(\frac{1}{q}-\frac{1}{p}\right), & \left(\frac{1}{p},\frac{1}{q}\right) \in \mathrm{T}_1. \end{cases} \tag{3.6}$$

Before describing the dilation property of the α-modulation spaces, we introduce some critical powers. Let us write $s_p = n(1/(1 \wedge p)-1)$ and

$$s_c = \begin{cases} R(p,q;1,\alpha), & \lambda > 1,\\ -R(p,q;\alpha,1), & \lambda \leqslant 1. \end{cases} \tag{3.7}$$


1/q
S_1
S_2
$\left(\frac{1}{2},\frac{1}{2}\right)$
S_3
(0,0)
(1,0)
1/p
1/q
(0, 1)
T_3
(1, 1)
T_2
$\left(\frac{1}{2},\frac{1}{2}\right)$
T_1
(0, 0)
1/p

Figure 1: Distribution of s_c. The left-hand side figure is for $\lambda > 1$, the right-hand side figure is for $\lambda \leqslant 1$.

Theorem 3.1. *Let $0 \leqslant \alpha < 1, \lambda > 0$ and $s + s_c \neq 0$. Then*

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim \lambda^{-\frac{n}{p}} \left[(1 \vee \lambda)^{s_p} \vee \lambda^{s+s_c}\right] \|f\|_{M^{s,\alpha}_{p,q}}. \tag{3.8}$$

holds for all $f \in M^{s,\alpha}_{p,q}$. Conversely, if

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim \lambda^{-\frac{n}{p}} F(\lambda) \|f\|_{M^{s,\alpha}_{p,q}}$$

holds for some $F : (0,\infty) \to (0,\infty)$ and all $f \in M^{s,\alpha}_{p,q}$, then $F(\lambda) \gtrsim (1 \vee \lambda)^{s_p} \vee \lambda^{s+s_c}$.

Proof. (**Sufficiency**) We denote by $\Box^\alpha_{k,1/\lambda}$ the pseudo-differential operator with symbol $(\eta^\alpha_k)_\lambda$. For every $l \in \mathbb{Z}^n$ and $\lambda > 0$, we introduce

$$\Lambda(l,\lambda) = \{k \in \mathbb{Z}^n : \Box^\alpha_{k,1/\lambda} \Box^\alpha_l f \neq 0\}. \tag{3.9}$$

For any $k \in \Lambda(l,\lambda)$, it follows from (1.1b) that k, l and λ satisfy

$$\lambda \langle l \rangle^{\frac{\alpha}{1-\alpha}} (l_j - C) < \langle k \rangle^{\frac{\alpha}{1-\alpha}} (k_j + C); \tag{3.10a}$$

$$\lambda \langle l \rangle^{\frac{\alpha}{1-\alpha}} (l_j + C) > \langle k \rangle^{\frac{\alpha}{1-\alpha}} (k_j - C) \tag{3.10b}$$

with $j = 1, 2, \cdots, n$. In view of (3.10), one sees that $k \in \Lambda(l,\lambda)$ is equivalent to $l \in \Lambda(k, 1/\lambda)$. Moreover, if (3.10) holds, then

$$\langle l \rangle \lesssim 1 \vee \lambda^{-(1-\alpha)} \quad \text{if and only if} \quad \langle k \rangle \lesssim 1 \vee \lambda^{1-\alpha}. \tag{3.11}$$

If $\langle l \rangle \gg 1 \vee \lambda^{-(1-\alpha)}$, without loss of generality, we may assume l belongs to some $\mathbb{R}^n_j$, when $l_j > 0$, from (3.10a); whereas when $l_j < 0$, from (3.10b)×(−1), we see $\langle k \rangle^{\frac{1}{1-\alpha}} \gtrsim \lambda \langle l \rangle^{\frac{1}{1-\alpha}}$. Conversely, for $k \in \Lambda(l,\lambda) \cap \mathbb{R}^n_j$, also from (3.10), we have $\langle k \rangle^{\frac{1}{1-\alpha}} \lesssim \lambda \langle l \rangle^{\frac{1}{1-\alpha}}$. Thus we have

$$\langle k \rangle \sim \lambda^{1-\alpha} \langle l \rangle. \tag{3.12}$$



Figure 2: 9 regions for the proof of Theorem 3.1

Since the volumes of $\mathrm{supp}(\eta_k^\alpha)_\lambda$ and $\mathrm{supp}\eta_l^\alpha$ are $O(\lambda^{-n}\langle k\rangle^{\frac{n\alpha}{1-\alpha}})$ and $O(\langle l\rangle^{\frac{n\alpha}{1-\alpha}})$, respectively, we see that

$$\#\Lambda(l,\lambda)\sim 1\vee\lambda^{n(1-\alpha)}. \tag{3.13}$$

When $q=1$, from Lemma 2.1, we have

$$\begin{aligned}\left\|\sum_{k\in\Gamma}\Box_k^\alpha f_\lambda\right\|_{M_{p,1}^{s,\alpha}}&=\sum_{k\in\Gamma}\langle k\rangle^{\frac{s}{1-\alpha}}\|\Box_k^\alpha f_\lambda\|_p\\&=\sum_{k\in\Gamma}\langle k\rangle^{\frac{s}{1-\alpha}}\|(\Box_{k,1/\lambda}^\alpha f)(\lambda\cdot)\|_p\\&=\lambda^{-\frac{n}{p}}\sum_{k\in\Gamma}\langle k\rangle^{\frac{s}{1-\alpha}}\|\Box_{k,1/\lambda}^\alpha f\|_p\\&\leqslant\lambda^{-\frac{n}{p}}\sum_{k\in\Gamma}\langle k\rangle^{\frac{s}{1-\alpha}}\sum_{l\in\Lambda(k,1/\lambda)}\|\Box_{k,1/\lambda}^\alpha\Box_l^\alpha f\|_p.\end{aligned} \tag{3.14}$$

Case 1. $\boldsymbol{\lambda\leqslant 1,(\frac{1}{p},\frac{1}{q})\in \mathrm{I}\cup\mathrm{II}}$. For $p=1,\infty$, we apply the same technique as that appeared in Proposition 2.3 to remove $\Box_{k,1/\lambda}^\alpha$ in (3.14). When $|k|\lesssim 1$, from (3.11) we see that for $l\in\Lambda(k,1/\lambda)$, there is $1\lesssim\langle l\rangle^{\frac{1}{1-\alpha}}\lesssim\frac{1}{\lambda}$, which leads to

$$\begin{aligned}\left\|\sum_{|k|\lesssim 1}\Box_k^\alpha f_\lambda\right\|_{M_{p,1}^{s,\alpha}}&\lesssim\lambda^{-\frac{n}{p}}\sum_{\langle l\rangle\lesssim\lambda^{-(1-\alpha)}}\|\Box_l^\alpha f\|_p\lesssim\lambda^{-\frac{n}{p}}\sum_{l\in\mathbb{Z}^n}(1\vee\lambda^s)\langle l\rangle^{\frac{s}{1-\alpha}}\|\Box_l^\alpha f\|_p\\&\lesssim\lambda^{-\frac{n}{p}}(1\vee\lambda^s)\|f\|_{M_{p,1}^{s,\alpha}}.\end{aligned} \tag{3.15}$$

By Plancherel's identity,

$$
\begin{aligned}
\left\|\sum_{|k|\lesssim 1}\Box_k^\alpha f_\lambda\right\|_{M_{2,2}^{s,\alpha}} &\lesssim \lambda^{-\frac{n}{2}}\left(\sum_{|k|\lesssim 1}\langle k\rangle^{\frac{2s}{1-\alpha}}\left\|\sum_{l\in\Lambda(k,1/\lambda)}\Box_{k,1/\lambda}^\alpha\Box_l^\alpha f\right\|_2^2\right)^{\frac{1}{2}}\\
&\lesssim \lambda^{-\frac{n}{2}}\left(\sum_{\langle l\rangle\lesssim\lambda^{-(1-\alpha)}}\|\Box_l^\alpha f\|_2^2\right)^{\frac{1}{2}}\lesssim \lambda^{-\frac{n}{2}}(1\vee\lambda^s)\|f\|_{M_{2,2}^{s,\alpha}}.
\end{aligned}
\tag{3.16}
$$

When $|k|\gg 1$, from (3.12)-(3.14), we see that

$$
\begin{aligned}
\left\|\sum_{|k|\gg 1}\Box_k^\alpha f_\lambda\right\|_{M_{p,1}^{s,\alpha}} &\lesssim \lambda^{-\frac{n}{p}+s}\sum_{|k|\gg 1}\sum_{l\in\Lambda(k,1/\lambda)}\langle l\rangle^{\frac{s}{1-\alpha}}\|\Box_l^\alpha f\|_p\\
&\lesssim \lambda^{-\frac{n}{p}+s}\sum_{l}\langle l\rangle^{\frac{s}{1-\alpha}}\sum_{k\in\Lambda(l,\lambda)}\|\Box_l^\alpha f\|_p\lesssim \lambda^{-\frac{n}{p}+s}\|f\|_{M_{p,1}^{s,\alpha}}.
\end{aligned}
\tag{3.17}
$$

In view of Plancherel's formula,

$$
\begin{aligned}
\left\|\sum_{|k|\gg 1}\Box_k^\alpha f_\lambda\right\|_{M_{2,2}^{s,\alpha}} &\leqslant \lambda^{-\frac{n}{2}+s}\left(\sum_{|k|\gg 1}\sum_{l\in\Lambda(k,1/\lambda)}\langle l\rangle^{\frac{2s}{1-\alpha}}\|\Box_{k,1/\lambda}^\alpha\Box_l^\alpha f\|_2^2\right)^{\frac{1}{2}}\\
&\lesssim \lambda^{-\frac{n}{2}+s}\left(\sum_{l}\langle l\rangle^{\frac{2s}{1-\alpha}}\sum_{k\in\Lambda(l,\lambda)}\|\Box_{k,1/\lambda}^\alpha\Box_l^\alpha f\|_2^2\right)^{\frac{1}{2}}\lesssim \lambda^{-\frac{n}{2}+s}\|f\|_{M_{2,2}^{s,\alpha}}.
\end{aligned}
\tag{3.18}
$$

Combining (3.15)-(3.18), we use complex interpolation to get

$$
\|f_\lambda\|_{M_{p,q}^{s,\alpha}}\lesssim\lambda^{-\frac{n}{p}}(1\vee\lambda^s)\|f\|_{M_{p,q}^{s,\alpha}}. \tag{3.19}
$$

Case 2. $\lambda>1$, $\left(\frac{1}{p},\frac{1}{q}\right)\in \mathrm{I}\cup\mathrm{III}^*$. Through the point $\left(\frac{1}{p},\frac{1}{q}\right)$, one can draw the parallel line to the $\frac{1}{q}$-axis. We assume there exists some $(\theta,\eta)\in[0,1]\times[0,1]$, such that the parallel line cuts the line segment connecting $(0,1),\left(\frac{1}{2},1\right)$ and the line segment connecting $(0,0),\left(\frac{1}{2},\frac{1}{2}\right)$ at $\left(\frac{\theta}{2},1\right)$ and $\left(\frac{\theta}{2},\frac{\theta}{2}\right)$, respectively. Assume that

$$
\frac{1}{p}=\frac{\theta}{2},\quad \frac{1}{q}=1-\left(1-\frac{\theta}{2}\right)\eta.
$$

When $|k|\lesssim\lambda^{1-\alpha}$, from (3.11) and (3.14), we have

$$
\left\|\sum_{|k|\lesssim\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\right\|_{M_{\infty,1}^{s,\alpha}}\leqslant\sum_{|l|\lesssim 1}\|\Box_l^\alpha f\|_\infty\sum_{k\in\Lambda(l,\lambda)}\langle k\rangle^{\frac{s}{1-\alpha}}\lesssim\left(1\vee\lambda^{s+n(1-\alpha)}\right)\|f\|_{M_{\infty,1}^{s,\alpha}}. \tag{3.20}
$$

By the Schwartz inequality and the Plancherel identity,

$$
\begin{aligned}
\left\|\sum_{|k|\lesssim\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\right\|_{M_{2,1}^{s,\alpha}} &\leqslant \lambda^{-\frac{n}{2}}\sum_{|l|\lesssim 1}\left(\sum_{k\in\Lambda(l,\lambda)}\|\Box_{k,1/\lambda}^\alpha\Box_l^\alpha f\|_2^2\right)^{\frac{1}{2}}\left(\sum_{k\in\Lambda(l,\lambda)}\langle k\rangle^{\frac{2s}{1-\alpha}}\right)^{\frac{1}{2}}\\
&\lesssim \lambda^{-\frac{n}{2}}\left(1\vee\lambda^{s+\frac{n}{2}(1-\alpha)}\right)\|f\|_{M_{2,1}^{s,\alpha}}.
\end{aligned}
\tag{3.21}
$$

From (3.11), we have

$$\begin{aligned}\Bigg\|\sum_{|k|\lesssim\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\Bigg\|_{M^{s,\alpha}_{\infty,\infty}} &\lesssim \sup_{|k|\lesssim\lambda^{1-\alpha}}\langle k\rangle^{\frac{s}{1-\alpha}}\|\Box_k^\alpha f_\lambda\|_\infty \\ &\lesssim (1\vee\lambda^s)\sup_{|k|\lesssim\lambda^{1-\alpha}}\sum_{l\in\Lambda(k,1/\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_\infty \\ &\lesssim (1\vee\lambda^s)\sup_{|l|\lesssim1}\|\Box_l^\alpha f\|_\infty\lesssim(1\vee\lambda^s)\|f\|_{M^{s,\alpha}_{\infty,\infty}}.\end{aligned}\tag{3.22}$$

In view of Plancherel's equality,

$$\begin{aligned}\Bigg\|\sum_{|k|\lesssim\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\Bigg\|_{M^{s,\alpha}_{2,2}} &\lesssim \left(\sum_{|k|\lesssim\lambda^{1-\alpha}}\langle k\rangle^{\frac{2s}{1-\alpha}}\|\Box_k^\alpha f_\lambda\|_2^2\right)^{\frac{1}{2}} \\ &\lesssim \lambda^{-\frac{n}{2}}\left(\sum_{|l|\lesssim1}(1\vee\lambda^s)^2\sum_{k\in\Lambda(l,\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_2^2\right)^{\frac{1}{2}} \\ &\lesssim \lambda^{-\frac{n}{2}}(1\vee\lambda^s)\left(\sum_{|l|\lesssim1}\|\Box_l^\alpha f\|_2^2\right)^{\frac{1}{2}}\lesssim\lambda^{-\frac{n}{2}}(1\vee\lambda^s)\|f\|_{M^{s,\alpha}_{2,2}}.\end{aligned}\tag{3.23}$$

When $|k|\gg\lambda^{1-\alpha}$, from (3.12)-(3.14), we have

$$\Bigg\|\sum_{|k|\gg\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\Bigg\|_{M^{s,\alpha}_{\infty,1}}\leqslant\lambda^s\sum_{|l|\gg1}\langle l\rangle^{\frac{s}{1-\alpha}}\sum_{k\in\Lambda(l,\lambda)}\|\Box_l^\alpha f\|_\infty\lesssim\lambda^{s+n(1-\alpha)}\|f\|_{M^{s,\alpha}_{\infty,1}}.\tag{3.24}$$

By Jensen's inequality,

$$\begin{aligned}\Bigg\|\sum_{|k|\gg\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\Bigg\|_{M^{s,\alpha}_{2,1}} &\lesssim \lambda^{-\frac{n}{2}+s}\sum_{|l|\gg1}\langle l\rangle^{\frac{s}{1-\alpha}}\sum_{k\in\Lambda(l,\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_2 \\ &\lesssim \lambda^{\frac{n}{2}+s}\sum_{|l|\gg1}\langle l\rangle^{\frac{s}{1-\alpha}}[\#\Lambda(l,\lambda)]^{\frac{1}{2}}\left(\sum_{k\in\Lambda(l,\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_2^2\right)^{\frac{1}{2}} \\ &\lesssim \lambda^{-\frac{n}{2}+s+\frac{n}{2}(1-\alpha)}\|f\|_{M^{s,\alpha}_{2,1}}.\end{aligned}\tag{3.25}$$

From (3.12),(3.13), we have

$$\begin{aligned}\Bigg\|\sum_{|k|\gg\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\Bigg\|_{M^{s,\alpha}_{\infty,\infty}} &\lesssim \sup_k\langle k\rangle^{\frac{s}{1-\alpha}}\|\Box^\alpha_{k,1/\lambda}f\|_\infty \\ &\lesssim \lambda^s\sup_k\sum_{l\in\Lambda(k,1/\lambda)}\langle l\rangle^{\frac{s}{1-\alpha}}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_\infty\lesssim\lambda^s\|f\|_{M^{s,\alpha}_{\infty,\infty}}.\end{aligned}\tag{3.26}$$

Similar to (3.18), one has that

$$\left\| \sum_{|k|\gg\lambda^{1-\alpha}} \Box_k^\alpha f_\lambda \right\|_{M_{2,2}^{s,\alpha}} \lesssim \lambda^{-\frac{n}{2}+s}\|f\|_{M_{2,2}^{s,\alpha}}. \tag{3.27}$$

Since $n(1-\alpha)\big(1-\frac{\theta}{2}\big) = (1-\theta)n(1-\alpha) + \theta\frac{n}{2}(1-\alpha)$, combining (3.20),(3.24),(3.21),(3.25), complex interpolation yields

$$\|f_\lambda\|_{M_{\frac{2}{\theta},1}^{s,\alpha}} \lesssim \lambda^{-\frac{\theta}{2}n}\left(1 \vee \lambda^{s+n(1-\alpha)\left(1-\frac{\theta}{2}\right)}\right)\|f\|_{M_{\frac{2}{\theta},1}^{s,\alpha}}. \tag{3.28}$$

Combining (3.22),(3.26),(3.23),(3.27), complex interpolation yields

$$\|f_\lambda\|_{M_{\frac{2}{\theta},\frac{2}{\theta}}^{s,\alpha}} \lesssim \lambda^{-\frac{\theta}{2}n}(1 \vee \lambda^s)\|f\|_{M_{\frac{2}{\theta},\frac{2}{\theta}}^{s,\alpha}}. \tag{3.29}$$

Interpolating (3.28) and (3.29), we have

$$\|f_\lambda\|_{M_{p,q}^{s,\alpha}} \lesssim \lambda^{-\frac{n}{p}}\left(1 \vee \lambda^{s+n(1-\alpha)\left(\frac{1}{q}-\frac{1}{p}\right)}\right)\|f\|_{M_{p,q}^{s,\alpha}}. \tag{3.30}$$

Case 3. $\boldsymbol{\lambda > 1, \big(\frac{1}{p}, \frac{1}{q}\big) \in \mathrm{II} \cup \mathrm{III}}$. Through the point $\big(\frac{1}{p}, \frac{1}{q}\big)$ one can make the parallel line to the $\frac{1}{q}$-axis. We assume there exists some $(\theta, \eta) \in [0,1]\times[0,1]$, such that the parallel line cuts the line segment connecting $(1,1), \big(\frac{1}{2}, 1\big)$ and the line segment connecting $(1,0), \big(\frac{1}{2}, \frac{1}{2}\big)$ at $\big(1-\frac{\theta}{2}, 1\big)$ and $\big(1-\frac{\theta}{2}, \frac{\theta}{2}\big)$, respectively. We can assume that

$$\frac{1}{p} = 1-\frac{\theta}{2}, \quad \frac{1}{q} = 1-\left(1-\frac{\theta}{2}\right)\eta.$$

When $|k| \lesssim \lambda^{1-\alpha}$, similarly to (3.20) and (3.22), we have

$$\left\| \sum_{|k|\lesssim\lambda^{1-\alpha}} \Box_k^\alpha f_\lambda \right\|_{M_{1,1}^{s,\alpha}} \lesssim \lambda^{-n}\left(1 \vee \lambda^{s+n(1-\alpha)}\right)\|f\|_{M_{1,1}^{s,\alpha}}; \tag{3.31}$$

$$\left\| \sum_{|k|\lesssim\lambda^{1-\alpha}} \Box_k^\alpha f_\lambda \right\|_{M_{1,\infty}^{s,\alpha}} \lesssim \lambda^{-n}\left(1 \vee \lambda^{s}\right)\|f\|_{M_{1,\infty}^{s,\alpha}}. \tag{3.32}$$

When $|k| \gg \lambda^{1-\alpha}$, similarly to (3.24) and (3.26), we have

$$\left\| \sum_{|k|\gg\lambda^{1-\alpha}} \Box_k^\alpha f_\lambda \right\|_{M_{1,1}^{s,\alpha}} \lesssim \lambda^{-n+s+n(1-\alpha)}\|f\|_{M_{1,1}^{s,\alpha}}; \tag{3.33}$$

$$\left\| \sum_{|k|\gg\lambda^{1-\alpha}} \Box_k^\alpha f_\lambda \right\|_{M_{1,\infty}^{s,\alpha}} \lesssim \lambda^{-n+s}\|f\|_{M_{1,\infty}^{s,\alpha}}. \tag{3.34}$$

Combining (3.31),(3.33),(3.21),(3.25), complex interpolation yields

$$\|f_\lambda\|_{M^{s,\alpha}_{\frac{2}{2-\theta},1}} \lesssim \lambda^{-\frac{n}{2}\left(1-\frac{\theta}{2}\right)} \left(1\vee \lambda^{s+n(1-\alpha)\left(1-\frac{\theta}{2}\right)}\right) \|f\|_{M^{s,\alpha}_{\frac{2}{2-\theta},1}}. \tag{3.35}$$

Combining (3.32),(3.34),(3.23),(3.27), complex interpolation yields

$$\|f_\lambda\|_{M^{s,\alpha}_{\frac{2}{2-\theta},\frac{2}{\theta}}} \lesssim \lambda^{-\frac{n}{2}\left(1-\frac{\theta}{2}\right)} \left(1\vee \lambda^{s}\right) \|f\|_{M^{s,\alpha}_{\frac{2}{2-\theta},\frac{2}{\theta}}}. \tag{3.36}$$

Interpolating (3.35) and (3.36), we have

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim \lambda^{-\frac{n}{p}} \left(1\vee \lambda^{s+n(1-\alpha)\left(\frac{1}{p}+\frac{1}{q}-1\right)}\right) \|f\|_{M^{s,\alpha}_{p,q}}. \tag{3.37}$$

Case 4. $\lambda \leqslant 1, \left(\frac{1}{p},\frac{1}{q}\right) \in \{\mathrm{I}^*\cup \mathrm{II}^*\cup \mathrm{III}\cup \mathrm{III}^*\}\backslash\{(0,1]\times[0,1]\}$; or $\lambda > 1, \left(\frac{1}{p},\frac{1}{q}\right) \in \mathrm{I}^*\cup \mathrm{II}^*\backslash\{(1,0)\}$. We observe that, for any $\left(\frac{1}{p},\frac{1}{q}\right)\in \mathrm{I}^*\cup\mathrm{II}^*$, we have $\left(\frac{1}{p^*},\frac{1}{q^*}\right)\in \mathrm{I}\cup\mathrm{II}$. By duality,

$$|\langle f_\lambda, g\rangle| = \frac{1}{\lambda^n}|\langle f, g_{\frac{1}{\lambda}}\rangle| \leqslant \frac{1}{\lambda^n}\|f\|_{M^{s,\alpha}_{p,q}}\left\|g_{\frac{1}{\lambda}}\right\|_{M^{-s,\alpha}_{p^*,q^*}}. \tag{3.38}$$

If we denote $\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim F(s,\lambda;p,q)\|f\|_{M^{s,\alpha}_{p,q}}$, from the previous several cases, we know that

$$\left\|g_{\frac{1}{\lambda}}\right\|_{M^{-s,\alpha}_{p^*,q^*}} \lesssim F\left(-s,\lambda^{-1};p^*,q^*\right)\|g\|_{M^{-s,\alpha}_{p^*,q^*}}. \tag{3.39}$$

By the principle of duality, it follows from (3.38) and (3.39) that

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim \lambda^{-n}F\left(-s,\lambda^{-1};p^*,q^*\right)\|f\|_{M^{s,\alpha}_{p,q}}. \tag{3.40}$$

For $\lambda\leqslant 1$ with $\left(\frac{1}{p},\frac{1}{q}\right)\in\{\mathrm{I}^*\cup\mathrm{III}\}\backslash\{1\}\times[0,1]$, from Case 2, (3.40) gives

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim \lambda^{-\frac{n}{p}} \left(1\vee \lambda^{s-n(1-\alpha)\left(\frac{1}{p}-\frac{1}{q}\right)}\right) \|f\|_{M^{s,\alpha}_{p,q}}. \tag{3.41}$$

For $\lambda\leqslant 1$ with $\left(\frac{1}{p},\frac{1}{q}\right)\in\{\mathrm{II}^*\cup\mathrm{III}^*\}\backslash\{(0,1)\}$, from Case 3, (3.40) gives

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim \lambda^{-\frac{n}{p}} \left(1\vee \lambda^{s-n(1-\alpha)\left(1-\frac{1}{q}-\frac{1}{p}\right)}\right) \|f\|_{M^{s,\alpha}_{p,q}}. \tag{3.42}$$

For $\lambda>1$ with $\left(\frac{1}{p},\frac{1}{q}\right)\in\mathrm{I}^*\cup\mathrm{II}^*\backslash\{(1,0)\}$, from Case 1, (3.40) gives

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim \lambda^{-\frac{n}{p}}(1\vee\lambda^s)\|f\|_{M^{s,\alpha}_{p,q}}. \tag{3.43}$$

Case 5. $\left(\frac{1}{p},\frac{1}{q}\right)\in\{\widehat{\mathrm{I}}\cup\widehat{\mathrm{II}}\}\backslash(1,\infty)\times(1,\infty)$. Since $\ell^q\subset\ell^1$, we know

$$\|\Box^\alpha_k f_\lambda\|_p \lesssim \lambda^{-\frac{n}{p}}\sum_{l\in\Lambda(k,1/\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_p \lesssim \lambda^{-\frac{n}{p}}\left(\sum_{l\in\Lambda(k,1/\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_p^q\right)^{\frac{1}{q}}. \tag{3.44}$$

From (3.44),(3.11), when $\lambda \leqslant 1$ and $|k| \lesssim 1$, we conclude that

$$\begin{aligned}\left\|\sum_{|k|\lesssim 1}\Box_k^\alpha f_\lambda\right\|_{M^{s,\alpha}_{\infty,q}} &\lesssim \left(\sum_{|k|\lesssim 1}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box_k^\alpha f_\lambda\|_\infty^q\right)^{\frac{1}{q}} \\ &\lesssim \left(\sum_{|k|\lesssim 1}\langle k\rangle^{\frac{sq}{1-\alpha}}\sum_{l\in\Lambda(k,1/\lambda)}\|\Box_{k,1/\lambda}^\alpha\Box_l^\alpha f\|_\infty^q\right)^{\frac{1}{q}} \\ &\lesssim \left(\sum_{\langle l\rangle\lesssim\lambda^{-(1-\alpha)}}\|\Box_l^\alpha f\|_\infty^q\right)^{\frac{1}{q}} \lesssim (1\vee\lambda^s)\|f\|_{M^{s,\alpha}_{\infty,q}};\end{aligned} \tag{3.45}$$

and when $\lambda > 1$ and $|k| \lesssim \lambda^{1-\alpha}$,

$$\begin{aligned}\left\|\sum_{|k|\lesssim\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\right\|_{M^{s,\alpha}_{\infty,q}} &\lesssim \left(\sum_{|k|\lesssim\lambda^{1-\alpha}}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box_k^\alpha f_\lambda\|_\infty^q\right)^{\frac{1}{q}} \\ &\lesssim \left(\sum_{|k|\lesssim\lambda^{1-\alpha}}\langle k\rangle^{\frac{sq}{1-\alpha}}\sum_{l\in\Lambda(k,1/\lambda)}\|\Box_l^\alpha f\|_\infty^q\right)^{\frac{1}{q}} \\ &\lesssim \left(\sum_{|l|\lesssim 1}\|\Box_l^\alpha f\|_\infty^q\sum_{|k|\lesssim\lambda^{1-\alpha}}\langle k\rangle^{\frac{sq}{1-\alpha}}\right)^{\frac{1}{q}} \\ &\lesssim \left(1\vee\lambda^{s+n(1-\alpha)\frac{1}{q}}\right)\|f\|_{M^{s,\alpha}_{\infty,q}}.\end{aligned} \tag{3.46}$$

When $|k| \gg 1\vee\lambda^{1-\alpha}$, from (3.44), (3.12), (3.13), we have

$$\begin{aligned}\left\|\sum_{|k|\gg 1\vee\lambda^{1-\alpha}}\Box_k^\alpha f_\lambda\right\|_{M^{s,\alpha}_{\infty,q}} &\lesssim \left(\sum_{|k|\gg 1\vee\lambda^{1-\alpha}}\langle k\rangle^{\frac{sq}{1-\alpha}}\sum_{l\in\Lambda(k,1/\lambda)}\|\Box_l^\alpha f\|_\infty^q\right)^{\frac{1}{q}} \\ &\lesssim \lambda^s\left(\sum_{|l|\gg 1}\langle l\rangle^{\frac{sq}{1-\alpha}}\sum_{k\in\Lambda(l,\lambda)}\|\Box_l^\alpha f\|_\infty\right)^{\frac{1}{q}} \\ &\lesssim \lambda^s(1\vee\lambda)^{n(1-\alpha)\frac{1}{q}}\|f\|_{M^{s,\alpha}_{\infty,q}}.\end{aligned} \tag{3.47}$$

Therefore, combining (3.45)-(3.47), we get

$$\|f_\lambda\|_{M^{s,\alpha}_{\infty,q}} \lesssim 1\vee\lambda^s\vee\lambda^{s+n(1-\alpha)\frac{1}{q}}\|f\|_{M^{s,\alpha}_{\infty,q}}. \tag{3.48}$$

The same for $M^{s,\alpha}_{1,q}$. Corresponding to (3.48), we get

$$\|f_\lambda\|_{M^{s,\alpha}_{1,q}} \lesssim \lambda^{-n}\left(1\vee\lambda^s\vee\lambda^{s+n(1-\alpha)\frac{1}{q}}\right)\|f\|_{M^{s,\alpha}_{1,q}}. \tag{3.49}$$

Whereas when $p=2, \lambda \leqslant 1$ and $|k| \lesssim 1$, from (3.44),(3.11), we have

$$\begin{aligned}
\Big\| \sum_{|k|\lesssim 1} \Box_k^\alpha f_\lambda \Big\|_{M_{2,q}^{s,\alpha}} &\lesssim \lambda^{-\frac{n}{2}} \left(\sum_{|k|\lesssim 1} \langle k\rangle^{\frac{sq}{1-\alpha}} \sum_{l\in\Lambda(k,1/\lambda)} \|\Box_{k,1/\lambda}^\alpha \Box_l^\lambda f\|_2^q \right)^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{2}} \left(\sum_{\langle l\rangle \lesssim \lambda^{-(1-\alpha)}} \|\Box_l^\alpha f\|_2^q \right)^{\frac{1}{q}} \lesssim \lambda^{-\frac{n}{2}} (1\vee \lambda^s) \|f\|_{M_{2,q}^{s,\alpha}}.
\end{aligned} \tag{3.50}$$

When $\lambda > 1, |k| \lesssim \lambda^{1-\alpha}$, by (3.44), (3.11) and Hölder's inequality, we have

$$\begin{aligned}
\Big\| \sum_{|k|\lesssim \lambda^{1-\alpha}} \Box_k^\alpha f_\lambda \Big\|_{M_{2,q}^{s,\alpha}} &\lesssim \lambda^{-\frac{n}{2}} \left(\sum_{|l|\lesssim 1} \sum_{k\in\Lambda(l,\lambda)} \langle k\rangle^{\frac{sq}{1-\alpha}} \|\Box_{k,1/\lambda}^\alpha \Box_l^\alpha f\|_2^q \right)^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{2}} \left[\sum_{|l|\lesssim 1} \left(\sum_{k\in\Lambda(l,\lambda)} \|\Box_{k,1/\lambda}^\alpha \Box_l^\alpha f\|_2^2 \right)^{\frac{q}{2}} \left(\sum_{|k|\lesssim\lambda^{1-\alpha}} \langle k\rangle^{\frac{sq}{1-\alpha}\cdot\frac{2}{2-q}} \right)^{\frac{2-q}{2}} \right]^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{2}} \left(1 \vee \lambda^{s+n(1-\alpha)\left(\frac{1}{q}-\frac{1}{2}\right)} \right) \|f\|_{M_{2,q}^{s,\alpha}}.
\end{aligned} \tag{3.51}$$

When $|k| \gg 1\vee\lambda^{1-\alpha}$, in view of (3.44),(3.12),(3.13) and Hölder's inequality, we have

$$\begin{aligned}
\Big\| \sum_{|k|\gg 1\vee\lambda^{1-\alpha}} \Box_k^\alpha f_\lambda \Big\|_{M_{2,q}^{s,\alpha}} &\lesssim \lambda^{-\frac{n}{2}+s} \left(\sum_{|l|\gg 1} \langle l\rangle^{\frac{sq}{1-\alpha}} \sum_{k\in\Lambda(l,\lambda)} \|\Box_l^\alpha \Box_{k,1/\lambda}^\alpha f\|_2^q \right)^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{2}+s} \left[\sum_{|l|\gg 1} \langle l\rangle^{\frac{sq}{1-\alpha}} [\#\Lambda(l,\lambda)]^{1-\frac{q}{2}} \left(\sum_{k\in\Lambda(l,\lambda)} \|\Box_l^\alpha \Box_{k,1/\lambda}^\alpha f\|_2^2 \right)^{\frac{q}{2}} \right]^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{2}+s} (1\vee\lambda)^{n(1-\alpha)\left(\frac{1}{q}-\frac{1}{2}\right)} \|f\|_{M_{2,q}^{s,\alpha}}.
\end{aligned} \tag{3.52}$$

Therefore, combining (3.50)-(3.52), we get

$$\|f_\lambda\|_{M_{2,q}^{s,\alpha}} \lesssim \lambda^{-\frac{n}{2}} \left(1\vee\lambda^s\vee\lambda^{s+n(1-\alpha)\left(\frac{1}{q}-\frac{1}{2}\right)} \right) \|f\|_{M_{2,q}^{s,\alpha}}. \tag{3.53}$$

For $\left(\frac{1}{p},\frac{1}{q}\right) \in \widehat{\mathrm{I}}$, complex interpolation between (3.48) and (3.53) yields

$$\|f_\lambda\|_{M_{p,q}^{s,\alpha}} \lesssim \lambda^{-\frac{n}{p}} \left(1\vee\lambda^s\vee\lambda^{s+n(1-\alpha)\left(\frac{1}{q}-\frac{1}{p}\right)} \right) \|f\|_{M_{p,q}^{s,\alpha}}; \tag{3.54}$$

while for $\left(\frac{1}{p},\frac{1}{q}\right) \in \widehat{\mathrm{II}}\backslash(1,\infty)\times(1,\infty)$, complex interpolation between (3.49) and (3.53) yields

$$\|f_\lambda\|_{M_{p,q}^{s,\alpha}} \lesssim \lambda^{-\frac{n}{p}} \left(1\vee\lambda^s\vee\lambda^{s+n(1-\alpha)\left(\frac{1}{p}+\frac{1}{q}-1\right)} \right) \|f\|_{M_{p,q}^{s,\alpha}}. \tag{3.55}$$

Case 6. $\boldsymbol{(\frac{1}{p},\frac{1}{q})\in\widehat{\mathbf{III}}}$**.** Since $l^p\subset l^1$, we know

$$\|\Box_k^\alpha f_\lambda\|_p=\lambda^{-\frac{n}{p}}\|\Box^\alpha_{k,1/\lambda}f\|_p\leqslant\lambda^{-\frac{n}{p}}\left(\sum_{l\in\Lambda(k,1/\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_p^p\right)^{\frac{1}{p}}. \tag{3.56}$$

When $\lambda\leqslant 1$ and $|k|\lesssim 1$, from (1.1b),(3.12), we see that

$$\text{diam supp}\mathscr{F}[(\mathscr{F}^{-1}(\eta^\alpha_k)_\lambda)(x-\cdot)\Box^\alpha_l f(\cdot)]\lesssim 1/\lambda,$$

By Proposition 2.1, imitating the processes as in (**??**)-(**??**), we get

$$\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_p\lesssim(1/\lambda)^{n\left(\frac{1}{p}-1\right)}\left(\lambda/\langle k\rangle^{\frac{\alpha}{1-\alpha}}\right)^{n\left(\frac{1}{p}-1\right)}\|\Box^\alpha_l f\|_p\lesssim\|\Box^\alpha_l f\|_p. \tag{3.57}$$

From (3.56), (3.57), the embedding $\ell^1\subset\ell^{\frac{q}{p}}$, and Hölder's inequality, we have

$$\begin{aligned}\left\|\sum_{|k|\lesssim 1}\Box^\alpha_k f_\lambda\right\|_{M^{s,\alpha}_{p,q}}&\lesssim\lambda^{-\frac{n}{p}}\left[\sum_{|k|\lesssim 1}\langle k\rangle^{\frac{sq}{1-\alpha}}\left(\sum_{l\in\Lambda(k,1/\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_p^p\right)^{\frac{q}{p}}\right]^{\frac{1}{q}}\\&\lesssim\lambda^{-\frac{n}{p}}\left(\sum_{|k|\lesssim 1}\langle k\rangle^{\frac{sp}{1-\alpha}}\sum_{l\in\Lambda(k,1/\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_p^p\right)^{\frac{1}{p}}\\&\lesssim\lambda^{-\frac{n}{p}}\left[\sum_{\langle l\rangle\lesssim\lambda^{-(1-\alpha)}}\left(\sum_{|k|\lesssim 1}\langle k\rangle^{\frac{s}{1-\alpha}\cdot\frac{pq}{q-p}}\right)^{\frac{q-p}{q}}\left(\sum_{|k|\lesssim 1}\|\Box^\alpha_l f\|_p^q\right)^{\frac{p}{q}}\right]^{\frac{1}{p}}\\&\lesssim\lambda^{-\frac{n}{p}}(1\vee\lambda^s)\|f\|_{M^{s,\alpha}_{p,q}}.\end{aligned} \tag{3.58}$$

When $\lambda\leqslant 1$ and $|k|\gg 1$, from (1.1b), (3.12), we see that

$$\text{diam supp}\mathscr{F}[(\mathscr{F}^{-1}(\eta^\alpha_k)_\lambda)(x-\cdot)\Box^\alpha_l f(\cdot)]\lesssim\langle k\rangle^{\frac{\alpha}{1-\alpha}}/\lambda.$$

Similarly to (3.57), we get

$$\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_p\lesssim\left(\langle k\rangle^{\frac{\alpha}{1-\alpha}}/\lambda\right)^{n\left(\frac{1}{p}-1\right)}\left(\lambda/\langle k\rangle^{\frac{\alpha}{1-\alpha}}\right)^{n\left(\frac{1}{p}-1\right)}\|\Box^\alpha_l f\|_p\leqslant\|\Box^\alpha_l f\|_p. \tag{3.59}$$

From (3.56), (3.59), (3.12), (3.13), we have

$$\begin{aligned}\left\|\sum_{|k|\gg 1}\Box^\alpha_k f_\lambda\right\|_{M^{s,\alpha}_{p,q}}&\lesssim\lambda^{-\frac{n}{p}}\left[\sum_{|k|\gg 1}\langle k\rangle^{\frac{sq}{1-\alpha}}\left(\sum_{l\in\Lambda(k,1/\lambda)}\|\Box^\alpha_{k,1/\lambda}\Box^\alpha_l f\|_p^p\right)^{\frac{q}{p}}\right]^{\frac{1}{q}}\\&\lesssim\lambda^{-\frac{n}{p}+s}\left(\sum_{|k|\gg 1}[\#\Lambda(k,1/\lambda)]^{\frac{q}{p}-1}\sum_{l\in\Lambda(k,1/\lambda)}\langle l\rangle^{\frac{sq}{1-\alpha}}\|\Box^\alpha_l f\|_p^q\right)^{\frac{1}{q}}\\&\lesssim\lambda^{-\frac{n}{p}+s-n(1-\alpha)\left(\frac{1}{p}-\frac{1}{q}\right)}\left(\sum_l\langle l\rangle^{\frac{sq}{1-\alpha}}\sum_{k\in\Lambda(l,\lambda)}\|\Box^\alpha_l f\|_p^q\right)^{\frac{1}{q}}\\&\lesssim\lambda^{-\frac{n}{p}+s-n(1-\alpha)\left(\frac{1}{p}-\frac{1}{q}\right)}\|f\|_{M^{s,\alpha}_{p,q}}.\end{aligned} \tag{3.60}$$

For $\lambda > 1$ and $|k| \lesssim \lambda^{1-\alpha}$, from (1.1b), (3.12), we know

$$\operatorname{diam} \operatorname{supp} \mathscr{F}[(\mathscr{F}^{-1}(\eta_k^\alpha)_\lambda)(x-\cdot)\Box_l^\alpha f(\cdot)] \lesssim 1.$$

Similarly to (3.57), we get

$$\|\Box_{k,1/\lambda}^\alpha \Box_l^\alpha f\|_p \lesssim \left(\lambda/\langle k\rangle^{\frac{\alpha}{1-\alpha}}\right)^{n\left(\frac{1}{p}-1\right)} \|\Box_l^\alpha f\|_p. \tag{3.61}$$

From (3.56), (3.61), (3.11), we have

$$\begin{aligned}
\left\|\sum_{|k|\lesssim\lambda^{1-\alpha}} \Box_k^\alpha f_\lambda\right\|_{M_{p,q}^{s,\alpha}} &\lesssim \lambda^{-\frac{n}{p}} \left[\sum_{|k|\lesssim\lambda^{1-\alpha}} \langle k\rangle^{\frac{sq}{1-\alpha}} \left(\sum_{l\in\Lambda(k,1/\lambda)} \|\Box_{k,1/\lambda}^\alpha \Box_l^\alpha f\|_p^p\right)^{\frac{q}{p}}\right]^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{p}+n\left(\frac{1}{p}-1\right)} \left[\sum_{|k|\lesssim\lambda^{1-\alpha}} \langle k\rangle^{\left[\frac{s}{1-\alpha}-\frac{n\alpha}{1-\alpha}\left(\frac{1}{p}-1\right)\right]q} \left(\sum_{|l|\lesssim 1} \|\Box_l^\alpha f\|_p^p\right)^{\frac{q}{p}}\right]^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{p}+n\left(\frac{1}{p}-1\right)} \left(1 \vee \lambda^{s-n\alpha\left(\frac{1}{p}-1\right)+n(1-\alpha)\frac{1}{q}}\right) \|f\|_{M_{p,q}^{s,\alpha}}.
\end{aligned} \tag{3.62}$$

When $\lambda > 1$ and $|k| \gg \lambda^{1-\alpha}$, similarly to (3.59), we get

$$\|\Box_{k,1/\lambda}^\alpha \Box_l^\alpha f\|_p \lesssim \|\Box_l^\alpha f\|_p. \tag{3.63}$$

From (3.56), (3.63), (3.12), and the embedding $\ell^1 \subset \ell^{\frac{q}{p}}$, we have

$$\begin{aligned}
\left\|\sum_{|k|\gg\lambda^{1-\alpha}} \Box_k^\alpha f_\lambda\right\|_{M_{p,q}^{s,\alpha}} &\lesssim \lambda^{-\frac{n}{p}} \left[\sum_{|k|\gg\lambda^{1-\alpha}} \langle k\rangle^{\frac{sq}{1-\alpha}} \left(\sum_{l\in\Lambda(k,1/\lambda)} \|\Box_{k,1/\lambda}^\alpha \Box_l^\alpha f\|_p^p\right)^{\frac{q}{p}}\right]^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{p}+s} \left(\sum_{|k|\gg\lambda^{1-\alpha}} \sum_{l\in\Lambda(k,1/\lambda)} \langle l\rangle^{\frac{sq}{1-\alpha}} \|\Box_l^\alpha f\|_p^q\right)^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{p}+s} \left(\sum_{|l|\gg 1} \langle l\rangle^{\frac{sq}{1-\alpha}} \sum_{k\in\Lambda(l,\lambda)} \|\Box_l^\alpha f\|_p^q\right)^{\frac{1}{q}} \\
&\lesssim \lambda^{-\frac{n}{p}+s+n(1-\alpha)\frac{1}{q}} \|f\|_{M_{p,q}^{s,\alpha}}.
\end{aligned} \tag{3.64}$$

We summarize the argument in this case as: if $\lambda \leqslant 1$, (3.58) and (3.60) give

$$\|f_\lambda\|_{M_{p,q}^{s,\alpha}} \lesssim \lambda^{-\frac{n}{p}} \left(1 \vee \lambda^{s+n(1-\alpha)\left(\frac{1}{q}-\frac{1}{p}\right)}\right) \|f\|_{M_{p,q}^{s,\alpha}}; \tag{3.65}$$

else if $\lambda > 1$, (3.62) and (3.64) give

$$\|f_\lambda\|_{M_{p,q}^{s,\alpha}} \lesssim \lambda^{-\frac{n}{p}+n\left(\frac{1}{p}-1\right)} \left(1 \vee \lambda^{s-n\alpha\left(\frac{1}{p}-1\right)+n(1-\alpha)\frac{1}{q}}\right) \|f\|_{M_{p,q}^{s,\alpha}}. \tag{3.66}$$

Case 7. $\boldsymbol{\left(\frac{1}{p}, \frac{1}{q}\right) \in \widehat{\mathbf{II}} \cap (1,\infty) \times (1,\infty)}$. It is a natural consequence of Cases 5 and 6 by complex interpolation.

(3.8) in the case $\lambda \geqslant 1$ follows from (3.30), (3.54), (3.37), (3.55), (3.43), (3.66). (3.8) in the case $\lambda < 1$ follows from (3.19), (3.54), (3.55), (3.41), (3.66), (3.42). □



Figure 3: The left-hand side figure is for $\alpha_1 \geqslant \alpha_2$; while the right-hand side figure is for $\alpha_1 < \alpha_2$.

Remark 3.1. *If $s = -s_c$, we have the substitution for* (3.8)*:*

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \lesssim \lambda^{-\frac{n}{p}} F(\lambda) \|f\|_{M^{s,\alpha}_{p,q}}, \tag{3.67}$$

where

$$F(\lambda) = \begin{cases} (\ln \lambda)^{0 \vee \left(\frac{1}{q} - \frac{1}{p}\right) \vee \left(\frac{1}{q} + \frac{1}{p} - 1\right)}, & \lambda > 1,\ p \geq 1; \\ \lambda^{n\left(\frac{1}{p} - 1\right)} (\ln \lambda)^{\frac{1}{q}}, & \lambda > 1,\ p \leq 1; \\ \left(\ln \frac{1}{\lambda}\right)^{0 \vee \left(\frac{1}{p} - \frac{1}{q}\right) \vee \left(1 - \frac{1}{p} - \frac{1}{q}\right)}, & \lambda \leqslant 1. \end{cases} \tag{3.68}$$

4 Embedding between α-modulation and Besov spaces

As $1 \leq p, q \leq \infty$, some sufficient conditions for the inclusions between modulation and Besov spaces were obtained by Gröbner [11], then Toft [16] improved Gröbner's sufficient conditions, which were proven to be necessary by Sugimoto and Tomita [15]. Their results were generalized to the cases $0 < p, q \leq \infty$ in [19, 20]. Gröbner [11] also considered the inclusions between α_1-modulation and α_2-modulation spaces for $1 \leq p, q \leq \infty$ and his results are optimal in the cases $(1/p, 1/q)$ is located in the vertices of the square $[0,1]^2$. We will improve Gröbner's results in the cases $1 \leq p, q \leq \infty$ and our results also cover the cases $0 < p < 1$ or $0 < q < 1$.

4.1 Embedding between α-modulation spaces

Theorem 4.1. *Let* $(\alpha_1, \alpha_2) \in [0, 1) \times [0, 1)$*. Then*

$$M_{p,q}^{s_1,\alpha_1} \subset M_{p,q}^{s_2,\alpha_2} \tag{4.1}$$

holds if and only if $s_1 \geqslant s_2 + R(p, q; \alpha_1, \alpha_2)$*.*

Remark. In the first versions of the paper, we obtain the sufficiency of Theorem 4.1, soon after Toft and Wahlberg [17] independently considered the embeddings between α-modulation and Besov spaces in the cases $1 \le p, q \le \infty$ and they first showed the necessity of Theorem 4.1 in the regions $(1/p, 1/q) \in (S_2 \cup S_3) \cap [0, 1]^2$ (see Fig. 3). After their work we can finally show the necessity of Theorem 4.1 in all cases.

Proof. (**Sufficiency**) For every $k \in \mathbb{Z}^n$, we introduce

$$\Lambda(k) = \{l \in \mathbb{Z}^n : \Box_l^{\alpha_2}\Box_k^{\alpha_1} f \neq 0\}. \tag{4.2}$$

If $\Box_l^{\alpha_2}\Box_k^{\alpha_1} f \neq 0$, then k and l satisfy

$$\begin{aligned} \langle l\rangle^{\frac{\alpha_2}{1-\alpha_2}}(l_j - C) &< \langle k\rangle^{\frac{\alpha_1}{1-\alpha_1}}(k_j + C), \\ \langle l\rangle^{\frac{\alpha_2}{1-\alpha_2}}(l_j + C) &> \langle k\rangle^{\frac{\alpha_1}{1-\alpha_1}}(k_j - C) \end{aligned} \tag{4.3}$$

for all $j = 1, 2, \cdots, n$. If $|k| \lesssim 1$, it is easy to see that $|l| \lesssim 1$. If $|k| \gg 1$, analogous to (3.12), we have

$$\langle l\rangle \sim \langle k\rangle^{\frac{1-\alpha_2}{1-\alpha_1}}. \tag{4.4}$$

Assume that $p \geqslant 1$, $q = 1$ and $s_2 = 0$, we have

$$\|f\|_{M_{p,1}^{0,\alpha_2}} \leqslant \sum_{k\in\mathbb{Z}^n} \sum_{l\in\Lambda(k)} \|\Box_k^{\alpha_1}\Box_l^{\alpha_2} f\|_p \lesssim \sum_k \#\Lambda(k)\|\Box_k^{\alpha_1} f\|_p. \tag{4.5}$$

We need an estimate of $\#\Lambda(k)$. Similar to (3.13), we have

$$\#\Lambda(k) \sim 1 \vee \langle k\rangle^{\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1}}. \tag{4.6}$$

If $p = 2$, inserting (4.6) into (4.5) and noticing (4.4), in view of Jensen's inequality we get

$$M_{2,1}^{s_2+0\vee\frac{n(\alpha_1-\alpha_2)}{2(1-\alpha_1)},\alpha_1} \hookrightarrow M_{2,1}^{s_2,\alpha_2}. \tag{4.7}$$

If $p = \infty$ or 1, from (4.5), (4.6), (4.4), one can directly obtain that

$$M_{\infty,1}^{s_2+0\vee\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1},\alpha_1} \hookrightarrow M_{\infty,1}^{s_2,\alpha_2}, \tag{4.8}$$

$$M_{1,1}^{s_2+0\vee\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1},\alpha_1} \hookrightarrow M_{1,1}^{s_2,\alpha_2}. \tag{4.9}$$

Case 1: $\left(\frac{1}{p}, \frac{1}{q}\right) \in \mathbf{I}$. For any $\theta \in [0, 1]$, $\left(\frac{\theta}{2}, 1\right)$ is at the line connecting $\left(\frac{1}{2}, 1\right)$ and $(0, 1)$. By complex interpolation between (4.7) and (4.8), one has that

$$M_{\frac{2}{\theta},1}^{0\vee\left(1-\frac{\theta}{2}\right)\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1},\alpha_1} \hookrightarrow M_{\frac{2}{\theta},1}^{0,\alpha_2}. \tag{4.10}$$

Since $\left(\frac{\theta}{2}, 1-\frac{\theta}{2}\right)$ is at the line segment by connecting $\left(\frac{1}{2}, \frac{1}{2}\right)$ and $(0,1)$, A complex interpolation combined with Proposition 2.7 and (4.8) yields

$$M^{0\vee(1-\theta)\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1}}_{\frac{2}{\theta},\frac{2}{2-\theta}} \hookrightarrow M^{0,\alpha_2}_{\frac{2}{\theta},\frac{2}{2-\theta}}. \tag{4.11}$$

For any $\left(\frac{1}{p}, \frac{1}{q}\right) \in \mathrm{I}$, we may suppose that there exists some $(\theta, \eta) \in [0,1] \times [0,1]$, such that

$$\frac{1}{p} = \frac{1}{p_1} = \frac{\theta}{2}, \qquad \frac{1}{q} = 1 - \frac{\theta\eta}{2}.$$

Therefore, a complex interpolation between (4.10) and (4.11) implies that

$$M^{s_2+\left(\frac{1}{q}-\frac{1}{p}\right)[0\vee n(\alpha_1-\alpha_2)],\alpha_1}_{p,q} \hookrightarrow M^{s_2,\alpha_2}_{p,q}. \tag{4.12}$$

Case 2: $\left(\frac{1}{p}, \frac{1}{q}\right) \in \mathbf{II}$. For any $\theta \in [0,1]$, $\left(1-\frac{\theta}{2}, 1-\frac{\theta}{2}\right)$ is at the line segment connecting $\left(\frac{1}{2}, \frac{1}{2}\right)$ and $(1,1)$. From Proposition 2.7 and (4.9), we see that

$$M^{s_2+(1-\theta)[0\vee n(\alpha_1-\alpha_2)],\alpha_1}_{\frac{2}{2-\theta},\frac{2}{2-\theta}} \hookrightarrow M^{s_2,\alpha_2}_{\frac{2}{2-\theta},\frac{2}{2-\theta}}. \tag{4.13}$$

Noticing that $\left(1-\frac{\theta}{2}, 1\right)$ is a point at the line segment connecting $\left(\frac{1}{2}, 1\right)$ and $(0,1)$, from (4.7) and (4.9), we see that

$$M^{s_2+\left(1-\frac{\theta}{2}\right)[0\vee n(\alpha_1-\alpha_2)],\alpha_1}_{\frac{2}{2-\theta},1} \hookrightarrow M^{s_2,\alpha_2}_{\frac{2}{2-\theta},1}. \tag{4.14}$$

Noticing that for any $\left(\frac{1}{p}, \frac{1}{q}\right) \in \mathrm{II}$, there exists some $(\theta, \eta) \in [0,1] \times [0,1]$ satisfying

$$\frac{1}{p} = 1 - \frac{\theta}{2}, \qquad \frac{1}{q} = 1 - \frac{\theta\eta}{2},$$

on the basis of (4.13) and (4.14), we conclude that

$$M^{s_2+\left(\frac{1}{p}+\frac{1}{q}-1\right)[0\vee n(\alpha_1-\alpha_2)],\alpha_1}_{p,q} \hookrightarrow M^{s_2,\alpha_2}_{p,q}. \tag{4.15}$$

When $\alpha_1 \leqslant \alpha_2$, (4.15) coincides with (4.12).
Case 3: $\left(\frac{1}{p}, \frac{1}{q}\right) \in \mathbf{I}^* \cup \mathbf{II}^*$. When $\left(\frac{1}{p}, \frac{1}{q}\right) \in \mathrm{I}^*$, $\left(\frac{1}{p^*}, \frac{1}{q^*}\right)$ is in I. From (4.12), we know

$$M^{-s_2,\alpha_2}_{p^*,q^*} \hookrightarrow M^{-s_2-\left(\frac{1}{p}-\frac{1}{q}\right)[0\vee n(\alpha_2-\alpha_1)],\alpha_1}_{p^*,q^*}.$$

The duality of α-modulation space implies that

$$M^{s_2+\left(\frac{1}{p}-\frac{1}{q}\right)[0\vee n(\alpha_2-\alpha_1)],\alpha_1}_{p,q} \hookrightarrow M^{s_2,\alpha_2}_{p,q}. \tag{4.16}$$

When $\left(\frac{1}{p}, \frac{1}{q}\right) \in \mathrm{II}^*$, by (4.15) and duality one has that

$$M^{s_2+\left(1-\frac{1}{p}-\frac{1}{q}\right)[0\vee n(\alpha_2-\alpha_1)],\alpha_1}_{p,q} \hookrightarrow M^{s_2,\alpha_2}_{p,q}. \tag{4.17}$$

When $\alpha_1 > \alpha_2$, (4.17) coincides with (4.16).

Case 4: $\left(\frac{1}{p},\frac{1}{q}\right)\in \mathrm{III}\cup\mathrm{III}^*$. We may assume that for any $\left(\frac{1}{p},\frac{1}{q}\right)\in$ III (III*), there exists a $\eta\in(0,1)$ satisfying

$$\frac{1}{q}=\eta\left(1-\frac{1}{p}\right)+\frac{1-\eta}{p}.$$

Notice that $\left(\frac{1}{p},\frac{1}{p}\right)$ and $\left(\frac{1}{p},1-\frac{1}{p}\right)$ are at the boundaries of II and I* (II* and I), respectively. If $\left(\frac{1}{p},\frac{1}{q}\right)\in$ III, a complex interpolation between (4.15) and (4.16) yields

$$M_{p,q}^{s_2+\left[n(\alpha_1-\alpha_2)\left(\frac{1}{p}+\frac{1}{q}-1\right)\vee n(\alpha_2-\alpha_1)\left(\frac{1}{p}-\frac{1}{q}\right)\right],\alpha_1}\hookrightarrow M_{p,q}^{s_2,\alpha_2}; \tag{4.18}$$

If $\left(\frac{1}{p},\frac{1}{q}\right)\in$ III*, a complex interpolation between (4.12) and (4.17) yields

$$M_{p,q}^{s_2+\left[n(\alpha_1-\alpha_2)\left(\frac{1}{q}-\frac{1}{p}\right)\vee n(\alpha_2-\alpha_1)\left(1-\frac{1}{p}-\frac{1}{q}\right)\right],\alpha_1}\hookrightarrow M_{p,q}^{s_2,\alpha_2}. \tag{4.19}$$

When $\alpha_1>\alpha_2$, (4.18) and (4.19) coincide with (4.15) and (4.12), respectively. When $\alpha_1\leqslant\alpha_2$, (4.18) and (4.19) coincide with (4.16) and (4.17), respectively.
Case 5: $\left(\frac{1}{p},\frac{1}{q}\right)\in\widehat{\mathrm{I}}$. Imitating the proof as in the counterpart of Theorem 3.1, we can easily get

$$M_{\infty,q}^{s_2+0\vee\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1}\frac{1}{q},\alpha_1}\subset M_{\infty,q}^{s_2,\alpha_2},\quad M_{2,q}^{s_2+0\vee\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1}\left(\frac{1}{q}-\frac{1}{2}\right),\alpha_1}\subset M_{2,q}^{s_2,\alpha_2}.$$

A complex interpolation yields

$$M_{p,q}^{s_2+0\vee\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1}\left(\frac{1}{q}-\frac{1}{p}\right),\alpha_1}\subset M_{p,q}^{s_2,\alpha_2}. \tag{4.20}$$

(4.20) coincides with (4.12).
Case 6: $\left(\frac{1}{p},\frac{1}{q}\right)\in\widehat{\mathrm{III}}$. From (1.1b), as well as (4.4), we see that

$$\operatorname{diam}\operatorname{supp}\mathscr{F}[\mathscr{F}^{-1}\eta_l^{\alpha_2}(x-\cdot)\Box_k^{\alpha_1}f(\cdot)]\lesssim\langle k\rangle^{\frac{\alpha_1\vee\alpha_2}{1-\alpha_1}},$$

In view of Proposition 2.1,

$$\|\Box_l^{\alpha_2}\Box_k^{\alpha_1}f\|_p\lesssim\langle k\rangle^{\left(\frac{1}{p}-1\right)\frac{0\vee n(\alpha_1-\alpha_2)}{1-\alpha_1}}\|\Box_k^{\alpha_1}f\|_p. \tag{4.21}$$

Inserting (4.21), (4.4), (4.6), from the embedding $\ell^p\subset\ell^1$ and with the aid of Jensen's inequality, we have

$$\begin{aligned}\|f\|_{M_{p,q}^{s_2,\alpha_2}}&\lesssim\left[\sum_{l\in\mathbb{Z}^n}\langle l\rangle^{\frac{s_2q}{1-\alpha_2}}\left(\sum_{k\in\Lambda(l)}\langle k\rangle^{(1-p)\frac{0\vee n(\alpha_1-\alpha_2)}{1-\alpha_1}}\|\Box_k^{\alpha_1}f\|_p^p\right)^{\frac{q}{p}}\right]^{\frac{1}{q}}\\&\lesssim\left(\sum_l\langle l\rangle^{\frac{s_2q}{1-\alpha_2}+\frac{0\vee n(\alpha_2-\alpha_1)}{1-\alpha_2}\left(\frac{q}{p}-1\right)}\sum_{k\in\Lambda(l)}\langle k\rangle^{nq\left(\frac{1}{p}-1\right)\frac{0\vee(\alpha_1-\alpha_2)}{1-\alpha_1}}\|\Box_k^{\alpha_1}f\|_p^q\right)^{\frac{1}{q}}\\&\lesssim\left(\sum_k\langle k\rangle^{\frac{s_2q}{1-\alpha_1}+\frac{0\vee n(\alpha_2-\alpha_1)}{1-\alpha_1}\left(\frac{q}{p}-1\right)+nq\frac{0\vee(\alpha_1-\alpha_2)}{1-\alpha_1}\left(\frac{1}{p}+\frac{1}{q}-1\right)}\|\Box_k^{\alpha_1}f\|_p^q\right)^{\frac{1}{q}}.\end{aligned} \tag{4.22}$$

When $\alpha_1 \leqslant \alpha_2$, (4.22) gives

$$M_{p,q}^{s_2+n(\alpha_2-\alpha_1)\left(\frac{1}{p}-\frac{1}{q}\right),\alpha_1} \subset M_{p,q}^{s_2,\alpha_2}; \tag{4.23}$$

whereas when $\alpha_1 > \alpha_2$, (4.22) gives

$$M_{p,q}^{s_2+n(\alpha_1-\alpha_2)\left(\frac{1}{p}+\frac{1}{q}-1\right),\alpha_1} \subset M_{p,q}^{s_2,\alpha_2}. \tag{4.24}$$

(4.23) and (4.24) coincide with (4.16) and (4.15), respectively.

Case 7: $\left(\boldsymbol{\frac{1}{p}}, \boldsymbol{\frac{1}{q}}\right) \in \widehat{\mathbf{II}}$**.** This is a consequence of the results in Cases 5 and 6 by complex interpolation. ☐

4.2 Embedding between Besov space and α-modulation space

In this section, we study the embedding between 1-modulation space and α-modulation spaces. In an analogous way to the previous subsection, we start with the embedding for the same indices p, q.

Theorem 4.2. *Let $\alpha \in [0,1)$. Then $B_{p,q}^{s_1} \subset M_{p,q}^{s_2,\alpha}$ holds if and only if $s_1 \geqslant s_2 + R(p,q;1,\alpha)$. Conversely, $M_{p,q}^{s_1,\alpha} \subset B_{p,q}^{s_2}$ holds if and only if $s_1 \geqslant s_2 + R(p,q;\alpha,1)$.*

Proof. (**sufficiency**) For every $j \in \mathbb{Z}_+$, we introduce

$$\Lambda(j) = \{k \in \mathbb{Z}^n : \Box_k^\alpha \triangle_j f \neq 0, \forall f \in \mathscr{S}'(\mathbb{R}^n)\}; \tag{4.25}$$

and for every $k \in \mathbb{Z}^n$, we introduce

$$\Lambda(k) = \{j \in \mathbb{Z}_+ : \Box_k^\alpha \triangle_j f \neq 0, \forall f \in \mathscr{S}'(\mathbb{R}^n)\}. \tag{4.26}$$

To a $j \in \mathbb{Z}_+$, for any $k \in \Lambda(j)$, it is easy to see that the quantitative relationship between k and j is

$$\langle k \rangle^{\frac{1}{1-\alpha}} \sim 2^j. \tag{4.27}$$

When $p \geqslant 1, q = 1$ and $s_2 = 0$, we have

$$\|f\|_{M_{p,1}^{0,\alpha}} \leqslant \sum_{k\in\mathbb{Z}^n} \sum_{j\in\Lambda(k)} \|\Box_k^\alpha \triangle_j f\|_p = \sum_{j\in\mathbb{Z}_+} \sum_{k\in\Lambda(j)} \|\triangle_j \Box_k^\alpha f\|_p. \tag{4.28}$$

For any $k \in \mathbb{Z}^n$ and any $j \in \mathbb{Z}_+$, it is easy to see

$$\#\Lambda(k) \sim 1, \quad \#\Lambda(j) \sim 2^{jn(1-\alpha)}. \tag{4.29}$$

Thus when $p = 2$, combining (4.28), (4.29), also with the aid of Jensen's inequality, we get

$$B_{2,1}^{s+\frac{n(1-\alpha)}{2}} \hookrightarrow M_{2,1}^{s,\alpha}. \tag{4.30}$$

If $p = 1$ or ∞, combining (4.29), (4.28), we get

$$B_{1,1}^{s+n(1-\alpha)} \hookrightarrow M_{1,1}^{s,\alpha}, \tag{4.31}$$

$$B_{\infty,1}^{s+n(1-\alpha)} \hookrightarrow M_{\infty,1}^{s,\alpha}. \tag{4.32}$$

Case1. $\left(\boldsymbol{\frac{1}{p}, \frac{1}{q}}\right) \in \mathbf{I}$. For any $\theta \in [0,1]$, a complex interpolation between (4.30) and (4.32) yields

$$B^{s+\left(1-\frac{\theta}{2}\right)n(1-\alpha)}_{\frac{2}{\theta},1} \hookrightarrow M^{s,\alpha}_{\frac{2}{\theta},1}; \tag{4.33}$$

while combined with Proposition 2.7 and (4.32), yields

$$B^{s+(1-\theta)n(1-\alpha)}_{\frac{2}{\theta},\frac{2}{2-\theta}} \hookrightarrow M^{s,\alpha}_{\frac{2}{\theta},\frac{2}{2-\theta}}. \tag{4.34}$$

In analogy to (4.12), we get from (4.33), (4.34) that

$$B^{s+n\left(\frac{1}{q}-\frac{1}{p}\right)(1-\alpha)}_{p,q} \hookrightarrow M^{s,\alpha}_{p,q}. \tag{4.35}$$

Conversely, when we encounter the embedding of α-modulation spaces into Besov spaces, for $2 \leqslant p \leqslant \infty$, considering (4.29), we have

$$\|f\|_{B^0_{p,1}} \leqslant \sum_{j\in\mathbb{Z}^+} \sum_{k\in\Lambda(j)} \|\square^\alpha_k \triangle_j f\|_p \lesssim \sum_{k\in\mathbb{Z}^n} \|\square^\alpha_k f\|_p = \|f\|_{M^{0,\alpha}_{p,1}}, \tag{4.36}$$

which gives

$$M^{s,\alpha}_{p,q} \hookrightarrow B^s_{p,q}. \tag{4.37}$$

Case2. $\left(\boldsymbol{\frac{1}{p}, \frac{1}{q}}\right) \in \mathbf{II}$. For any $\theta \in [0,1]$, a complex interpolation between (4.30) and (4.31) yields

$$B^{s+\left(1-\frac{\theta}{2}\right)n(1-\alpha)}_{\frac{2}{2-\theta},1} \hookrightarrow M^{s,\alpha}_{\frac{2}{2-\theta},1}. \tag{4.38}$$

From Proposition 2.7 and (4.31) it follows that

$$B^{s+(1-\theta)n(1-\alpha)}_{\frac{2}{2-\theta},\frac{2}{2-\theta}} \hookrightarrow M^{s,\alpha}_{\frac{2}{2-\theta},\frac{2}{2-\theta}}. \tag{4.39}$$

Analogous to (4.15), one can conclude from (4.38) and (4.39) that

$$B^{s+n\left(\frac{1}{q}+\frac{1}{p}-1\right)(1-\alpha)}_{p,q} \hookrightarrow M^{s,\alpha}_{p,q}. \tag{4.40}$$

Considering the embedding of α-modulation spaces into Besov spaces, (4.37) still holds if $(1/p, 1/q) \in II$.

Case3. $\left(\boldsymbol{\frac{1}{p}, \frac{1}{q}}\right) \in \mathbf{I^*} \cup \mathbf{II^*}$. Since $\left(\frac{1}{p^*}, \frac{1}{q^*}\right) \in \mathrm{I} \cup \mathrm{II}$, from (4.37), we see that $M^{-s,\alpha}_{p^*,q^*} \hookrightarrow B^{-s}_{p^*,q^*}$. Thus, the duality between $B^s_{p,q}$ and $B^{-s}_{p^*,q^*}$, as well as between $M^{s,\alpha}_{p,q}$ and $M^{-s,\alpha}_{p^*,q^*}$, implies that

$$B^s_{p,q} \hookrightarrow M^{s,\alpha}_{p,q}. \tag{4.41}$$

Conversely, if one considers the embedding of α-modulation space into Besov space, it follows from Theorem 2.1 and (4.35) that

$$M^{s+n(\alpha-1)\left(\frac{1}{q}-\frac{1}{p}\right),\alpha}_{p,q} \subset B^s_{p,q}, \quad \left(\frac{1}{p}, \frac{1}{q}\right) \in \mathrm{I}^*; \tag{4.42}$$

and from (4.40) it follows that

$$M^{s+n(\alpha-1)\left(\frac{1}{p}+\frac{1}{q}-1\right),\alpha}_{p,q} \subset B^s_{p,q}, \quad \left(\frac{1}{p}, \frac{1}{q}\right) \in \mathrm{II}^*. \tag{4.43}$$

Case4. $\boldsymbol{\left(\frac{1}{p},\frac{1}{q}\right)\in \mathrm{III}\cup \mathrm{III}^*}$. If $\left(\frac{1}{p},\frac{1}{q}\right)\in \mathrm{III}^*$, (4.35) and (4.41) contain that

$$B^{s+n\left(1-\frac{2}{p}\right)(1-\alpha)}_{p,\frac{p}{p-1}} \hookrightarrow M^{s,\alpha}_{p,\frac{p}{p-1}}, \quad B^{s}_{p,p} \hookrightarrow M^{s,\alpha}_{p,p}.$$

By interpolation we have

$$B^{s+n\left(\frac{1}{q}-\frac{1}{p}\right)(1-\alpha)}_{p,q} \hookrightarrow M^{s,\alpha}_{p,q}, \tag{4.44}$$

which coincides with (4.35). If $\left(\frac{1}{p},\frac{1}{q}\right)\in \mathrm{III}$, (4.40) and (4.41) imply that

$$B^{s}_{p,\frac{p}{p-1}} \subset M^{s,\alpha}_{p,\frac{p}{p-1}}, \quad B^{s+n\left(\frac{2}{p}-1\right)(1-\alpha)}_{p,p} \subset M^{s,\alpha}_{p,q}.$$

By interpolation,

$$B^{s+n\left(\frac{1}{q}+\frac{1}{p}-1\right)(1-\alpha)}_{p,q} \hookrightarrow M^{s,\alpha}_{p,q}, \tag{4.45}$$

which coincides with (4.40).

Conversely, considering the embedding of α-modulation space into Besov space, in view of Theorem 2.1 and (4.44) we have

$$M^{s+n(\alpha-1)\left(\frac{1}{q}-\frac{1}{p}\right),\alpha}_{p,q} \subset B^{s}_{p,q}, \quad \left(\frac{1}{p},\frac{1}{q}\right)\in \mathrm{III}; \tag{4.46}$$

while for $\left(\frac{1}{p},\frac{1}{q}\right)\in \mathrm{III}^*$, from (4.45), we have

$$M^{s+n(\alpha-1)\left(\frac{1}{p}+\frac{1}{q}-1\right),\alpha}_{p,q} \subset B^{s}_{p,q}. \tag{4.47}$$

(4.46) and (4.47) coincide with (4.42) and (4.43), respectively.

Case5 :$\boldsymbol{\left(\frac{1}{p},\frac{1}{q}\right)\in \widehat{\mathrm{I}}}$**.** For the embedding of Besov space into α-modulation space, imitating the argument in Theorem 4.1, we get

$$M^{s+n(1-\alpha)\frac{1}{q},1}_{\infty,q} \subset M^{s,\alpha}_{p,q}, \quad M^{s+n(1-\alpha)\left(\frac{1}{q}-\frac{1}{2}\right),1}_{2,q} \subset M^{s,\alpha}_{p,q}.$$

From them, we interpolate out

$$M^{s+n(1-\alpha)\left(\frac{1}{q}-\frac{1}{p}\right),1}_{p,q} \subset M^{s,\alpha}_{p,q}, \tag{4.48}$$

which coincides with (4.35). Conversely for the embedding of α-modulation space into Besov space, we have

$$M^{s,\alpha}_{p,q} \subset B^{s}_{p,q}, \tag{4.49}$$

which coincides with (4.37).

Case 6. $\boldsymbol{\left(\frac{1}{p},\frac{1}{q}\right)\in \widehat{\mathrm{III}}}$**.** If $\Box^{\alpha}_{k}\triangle_j \neq 0$, then

$$\mathrm{diam\ supp}\mathscr{F}[\mathscr{F}^{-1}\eta^{\alpha}_{k}(x-\cdot)\triangle_j f(\cdot)] \lesssim \langle k\rangle^{\frac{1}{1-\alpha}} \sim 2^{j}, \tag{4.50}$$

So, in view of Proposition 2.1 we have

$$\|\Box^{\alpha}_{k}\triangle_j f\|_p \lesssim \langle k\rangle^{\frac{n}{1-\alpha}\left(\frac{1}{p}-1\right)}\langle k\rangle^{-\frac{n\alpha}{1-\alpha}\left(\frac{1}{p}-1\right)}\|\triangle_j f\|_p \sim 2^{jn(1-\alpha)\left(\frac{1}{p}-1\right)}\|\triangle_j f\|_p. \tag{4.51}$$

From (4.51), (4.29), $\ell^p \subset \ell^1$ and Jensen's inequality it follows that

$$\begin{aligned}
\|f\|_{M^{s,\alpha}_{p,q}} &\lesssim \left[\sum_{k\in\mathbb{Z}^n} \langle k\rangle^{\frac{sq}{1-\alpha}} \left(\sum_{j\in\Lambda(k)} \|\Box^\alpha_k \triangle_j f\|^p_p\right)^{\frac{q}{p}}\right]^{\frac{1}{q}} \\
&\lesssim \left(\sum_k \langle k\rangle^{\left[\frac{s}{1-\alpha}+n\left(\frac{1}{p}-1\right)\right]q} \sum_{j\in\Lambda(k)} \|\triangle_j f\|^q_p\right)^{\frac{1}{q}} \qquad (4.52)\\
&\lesssim \left(\sum_j 2^{j\left[s+n(1-\alpha)\left(\frac{1}{p}-1\right)\right]q} \sum_{k\in\Lambda(j)} \|\triangle_j f\|^q_p\right)^{\frac{1}{q}} \lesssim \|f\|_{B^{s+n(1-\alpha)\left(\frac{1}{p}+\frac{1}{q}-1\right)}_{p,q}},
\end{aligned}$$

which implies that

$$B^{s+n(1-\alpha)\left(\frac{1}{p}+\frac{1}{q}-1\right),\alpha}_{p,q} \subset M^{s,\alpha}_{p,q}. \qquad (4.53)$$

It coincides with (4.40).

Conversely, when we study the embedding of α-modulation space into Besov space, in analogy to (4.50), we see

$$\operatorname{diam}\operatorname{supp}\mathscr{F}[\mathscr{F}^{-1}\varphi_j(x-\cdot)\Box^\alpha_k f(\cdot)] \lesssim 2^j$$

In contrast to (4.51), we conclude

$$\|\triangle_j \Box^\alpha_k f\|_p \lesssim 2^{jn\left(\frac{1}{p}-1\right)} 2^{-jn\left(\frac{1}{p}-1\right)} \|\Box^\alpha_k f\|_p = \|\Box^\alpha_k f\|_p. \qquad (4.54)$$

Inserting (4.54), the substitution for (4.52) is

$$\begin{aligned}
\|f\|_{B^s_{p,q}} &\lesssim \left[\sum_{j\in\mathbb{Z}^+} 2^{jsq} \left(\sum_{j\in\Lambda(j)} \|\triangle_j \Box^\alpha_k f\|^p_p\right)^{\frac{q}{p}}\right]^{\frac{1}{q}} \\
&\lesssim \left(\sum_j 2^{jsq} \#\Lambda(j)^{\frac{q}{p}-1} \sum_{k\in\Lambda(j)} \|\triangle_j \Box^\alpha_k f\|^q_p\right)^{\frac{1}{q}} \\
&\lesssim \left(\sum_k \langle k\rangle^{\left[s+n(1-\alpha)\left(\frac{1}{p}-\frac{1}{q}\right)\right]q} \|\Box^\alpha_k f\|^q_p\right)^{\frac{1}{q}} \lesssim \|f\|_{M^{s+n(1-\alpha)\left(\frac{1}{p}-\frac{1}{q}\right)}_{p,q}},
\end{aligned}$$

which implies that

$$M^{s+n(\alpha-1)\left(\frac{1}{q}-\frac{1}{p}\right),\alpha}_{p,q} \subset M^{s,1}_{p,q}. \qquad (4.55)$$

It coincides with (4.42).

Case7 $:\left(\boldsymbol{\frac{1}{p}}, \boldsymbol{\frac{1}{q}}\right) \in \widehat{\mathbf{II}}$. It is interpolated out from Case 5 and Case 6. $\Box$



Figure 4: Distribution of s_0

5 Multiplication algebra

It is well known that $B^s_{p,q}$ is a multiplication algebra if $s > n/p$, cf. [18]. But for α-modulation space, this issue is much more complicated. The regularity indices for which $M^{s,\alpha}_{p,q}$ constitutes a multiplication algebra, are quite different from those of Besov and modulation spaces.

We introduce a parameter, denoted by $s_0 = s_0(p, q; \alpha)$, to describe the regularity for which $M^{s,\alpha}_{p,q}$ with $s > s_0$ forms a multiplication algebra. Denote (see Figure 4)

$$\mathrm{D}_1 = \left\{ \left(\tfrac{1}{p}, \tfrac{1}{q}\right) \in \mathbb{R}^2_+ : \tfrac{1}{q} \geqslant \tfrac{2}{p}, \tfrac{1}{p} \leqslant \tfrac{1}{2} \right\}, \quad D_2 = \mathbb{R}^2_+ \setminus D_1$$

and

$$s_0 = \begin{cases} \frac{n\alpha}{p} + n(1-\alpha)\big(1 - 1 \wedge \frac{1}{q}\big) + \frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{q} - \frac{2}{p}\big), & \big(\frac{1}{p}, \frac{1}{q}\big) \in \mathrm{D}_1; \\ \frac{n\alpha}{p} + n(1-\alpha)\big(1 \vee \frac{1}{p} \vee \frac{1}{q} - \frac{1}{q}\big) + \frac{n\alpha(1-\alpha)}{2-\alpha}\big(1 \vee \frac{1}{p} \vee \frac{1}{q} - 1\big), & \big(\frac{1}{p}, \frac{1}{q}\big) \in \mathrm{D}_2. \end{cases}$$

Theorem 5.1. *If $s > s_0$, then $M^{s,\alpha}_{p,q}$ is a multiplication algebra, which is equivalent to say that for any $f, g \in M^{s,\alpha}_{p,q}$, we have*

$$\|fg\|_{M^{s,\alpha}_{p,q}} \lesssim \|f\|_{M^{s,\alpha}_{p,q}} \|g\|_{M^{s,\alpha}_{p,q}}. \tag{5.1}$$

In Section 7 we will give some counterexamples to show that s_0 is sharp if $(1/p, 1/q) \in \mathrm{D}_2 \cap \{p \geq 1\}$. When $\big(1/p,\ 1/q\big) \in D_1$, it is not very clear for us to know the sharp low bound of the index s for which $M^{s,\alpha}_{p,q}$ constitutes a multiplication algebra. As a straightforward consequence of Theorem 5.1, we have the following result for which $M^s_{p,q}$ is an algebra.

Corrolary 5.1. *Assume that*

$$s > \begin{cases} n\big(1 - 1 \wedge \frac{1}{q}\big), & \big(\frac{1}{p}, \frac{1}{q}\big) \in D_1; \\ n\big(1 \vee \frac{1}{p} \vee \frac{1}{q} - \frac{1}{q}\big), & \big(\frac{1}{p}, \frac{1}{q}\big) \in \mathbb{R}^2_+ \setminus D_1. \end{cases}$$

Then $M^s_{p,q}$ is a multiplication algebra, i.e.,

$$\|fg\|_{M^s_{p,q}} \lesssim \|f\|_{M^s_{p,q}} \|g\|_{M^s_{p,q}} \tag{5.2}$$

holds for all $f, g \in M^s_{p,q}$.

A natural long standing question on modulation, α-modulation and Besov spaces is: Can we reformulate α-modulation spaces by interpolations between modulation and Besov spaces, say,

$$M^{s,\alpha}_{p,q} = (M^{s_0,0}_{p,q},\ M^{s_1,1}_{p,q})_\alpha, \quad \text{if} \ \ s = (1-\alpha)s_0 + \alpha s_1? \tag{5.3}$$

The answer is negative at least for some special cases. Indeed, we see $M^{s_1,1}_{p,q}$ and $M^{s_0,0}_{p,q}$ are algebra if $s_1 > n/p$ and $s_0 > 0$. If (5.3) holds, then $M^{s,\alpha}_{p,q}$ is an algebra if $s > n\alpha/p$, however, this is not true if $1 < p < 2,\ 0 < q < 1$, see Section 7.

Corrolary 5.2. *Let $0 < \alpha < 1$. Then (5.3) does not hold if $1 < p < 2,\ 0 < q < 1$ and $s_0 = 0_+,\ s_1 = n/p$.*

Proof of Theorem 5.1. We start with some notations and basic conclusions. For every $(k^{(1)}, k^{(2)}) \in \mathbb{Z}^{2n}$, we introduce

$$\Lambda\big(k^{(1)}, k^{(2)}\big) = \big\{k \in \mathbb{Z}^n : \Box^\alpha_k (\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g) \neq 0\big\}; \tag{5.4}$$

and for every $k \in \mathbb{Z}^n$, we introduce

$$\Lambda(k) = \Big\{\big(k^{(1)}, k^{(2)}\big) \in \mathbb{Z}^{2n} : \Box^\alpha_k (\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g) \neq 0\Big\}.$$

It is worth to mention that $\Lambda\big(k^{(1)}, k^{(2)}\big)$ and $\Lambda(k)$ are independent of f and g. From (1.1b) we see that for any $k \in \Lambda(k^{(1)}, k^{(2)})$, or $(k^{(1)}, k^{(2)}) \in \Lambda(k)$, $k^{(1)}, k^{(2)}$ and k satisfy

$$\langle k\rangle^{\frac{\alpha}{1-\alpha}} (k_j - C) < \langle k^{(1)}\rangle^{\frac{\alpha}{1-\alpha}} (k^{(1)}_j + C) + \langle k^{(2)}\rangle^{\frac{\alpha}{1-\alpha}} (k^{(2)}_j + C), \tag{5.5a}$$

$$\langle k\rangle^{\frac{\alpha}{1-\alpha}} (k_j + C) > \langle k^{(1)}\rangle^{\frac{\alpha}{1-\alpha}} (k^{(1)}_j - C) + \langle k^{(2)}\rangle^{\frac{\alpha}{1-\alpha}} (k^{(2)}_j - C) \tag{5.5b}$$

for $j = 1, 2, \cdots, n$. Let $\langle k_{\max}\rangle$, $\langle k_{\min}\rangle$ and $\langle k_{\mathrm{med}}\rangle$ to be the maximal, minimal and medial ones in $\langle k\rangle, \langle k^{(1)}\rangle$ and $\langle k^{(2)}\rangle$, respectively. If (5.5) holds, then

$$\langle k_{\max}\rangle \sim \langle k_{\mathrm{med}}\rangle, \quad \#\big\{k_{\max} :\ k_{\max},\ k_{\mathrm{med}},\ k_{\min} \text{ satisfy (5.5)}\big\} \lesssim 1. \tag{5.6}$$

We write

$$\begin{aligned} K_j\big(k^{(1)}, k^{(2)}\big) &= \langle k^{(1)}\rangle^{\frac{\alpha}{1-\alpha}} k^{(1)}_j + \langle k^{(2)}\rangle^{\frac{\alpha}{1-\alpha}} k^{(2)}_j; \\ K\big(k^{(1)}, k^{(2)}\big) &= \max_{1\leqslant j\leqslant n} |K_j(k^{(1)}, k^{(2)})|. \end{aligned} \tag{5.7}$$

In order to get more precise estimates, we divide $\mathbb{Z}^{2n}$ of all $(k^{(1)}, k^{(2)})$ into

$$\begin{aligned} \Omega_0 &= \Big\{\big(k^{(1)}, k^{(2)}\big) \in \mathbb{Z}^{2n} : \langle k^{(1)}\rangle \sim \langle k^{(2)}\rangle\Big\}, \\ \Omega_1 &= \Big\{\big(k^{(1)}, k^{(2)}\big) \in \mathbb{Z}^{2n} : \langle k^{(1)}\rangle \gg \langle k^{(2)}\rangle\Big\}, \\ \Omega_2 &= \Big\{\big(k^{(1)}, k^{(2)}\big) \in \mathbb{Z}^{2n} : \langle k^{(1)}\rangle \ll \langle k^{(2)}\rangle\Big\}, \end{aligned} \tag{5.8}$$

and separate Ω_0 into

$$\begin{aligned}
\Omega_{0,1} &= \Big\{ \big(k^{(1)}, k^{(2)}\big) \in \Omega_0 : K\big(k^{(1)}, k^{(2)}\big) \lesssim \langle k^{(1)} \rangle^{\frac{\alpha}{1-\alpha}} \Big\};\\
\Omega_{0,2} &= \Big\{ \big(k^{(1)}, k^{(2)}\big) \in \Omega_0 : K\big(k^{(1)}, k^{(2)}\big) \gg \langle k^{(1)} \rangle^{\frac{\alpha}{1-\alpha}} \Big\}.
\end{aligned} \tag{5.9}$$

If $(k^{(1)}, k^{(2)}) \in \Omega_{0,1}$, from (5.5) it is easy to see that

$$\langle k \rangle \lesssim \langle k^{(1)} \rangle^{\alpha}. \tag{5.10}$$

Let $(k^{(1)}, k^{(2)}) \in \Omega_{0,2}$ be fixed. There exists some $y := y(k^{(1)}, k^{(2)}) \in (\alpha, 1]$ such that

$$K_i(k^{(1)}, k^{(2)}) = K(k^{(1)}, k^{(2)}) \sim \langle k^{(1)} \rangle^{\frac{y}{1-\alpha}}, \tag{5.11}$$

for some i with $1 \leqslant i \leqslant n$. We can assume that $K_i(k^{(1)}, k^{(2)}) > 0$. By (5.5) and (5.11) we have

$$\langle k \rangle^{\frac{\alpha}{1-\alpha}} (k_i - C) < K_i(k^{(1)}, k^{(2)}) + C(\langle k^{(1)} \rangle^{\frac{\alpha}{1-\alpha}} + \langle k^{(2)} \rangle^{\frac{\alpha}{1-\alpha}}) \lesssim \langle k^{(1)} \rangle^{\frac{y}{1-\alpha}}, \tag{5.12a}$$

$$\langle k \rangle^{\frac{\alpha}{1-\alpha}} (k_i + C) > K_i(k^{(1)}, k^{(2)}) - C(\langle k^{(1)} \rangle^{\frac{\alpha}{1-\alpha}} + \langle k^{(2)} \rangle^{\frac{\alpha}{1-\alpha}}) \gtrsim \langle k^{(1)} \rangle^{\frac{y}{1-\alpha}}. \tag{5.12b}$$

For every $k \in \Lambda(k^{(1)}, k^{(2)}) \cap \mathbb{R}^n_{\widetilde{i}}$, we substitute $\widetilde{i}$ for j in (5.5), thus (5.5a) and (5.5b) are rewritten as

$$\langle k \rangle^{\frac{\alpha}{1-\alpha}} (k_{\widetilde{i}} - C) < K_{\widetilde{i}}(k^{(1)}, k^{(2)}) + C(\langle k^{(1)} \rangle^{\frac{\alpha}{1-\alpha}} + \langle k^{(2)} \rangle^{\frac{\alpha}{1-\alpha}}); \tag{5.13a}$$

$$\langle k \rangle^{\frac{\alpha}{1-\alpha}} (k_{\widetilde{i}} + C) > K_{\widetilde{i}}(k^{(1)}, k^{(2)}) - C(\langle k^{(1)} \rangle^{\frac{\alpha}{1-\alpha}} + \langle k^{(2)} \rangle^{\frac{\alpha}{1-\alpha}}). \tag{5.13b}$$

For such k, we claim that

$$\langle k \rangle \sim \langle k^{(1)} \rangle^{y}. \tag{5.14}$$

(5.14) is obvious when $|k^{(1)}| \lesssim 1$ and so, it suffices to consider (5.14) in the case $|k^{(1)}| \gg 1$. If either $|k_i| \sim \langle k \rangle$ or $|K_{\widetilde{i}}(k^{(1)}, k^{(2)})| \sim K(k^{(1)}, k^{(2)})$ exists, (5.14) follows from (5.12) or (5.13) directly. Otherwise, we see that $|k_i| \ll \langle k \rangle$ and $|K_{\widetilde{i}}(k^{(1)}, k^{(2)})| \ll K(k^{(1)}, k^{(2)})$. When $k_{\widetilde{i}} > 0$, we let (5.13a)+(5.12a) and (5.13b)+(5.12b); whereas when $k_{\widetilde{i}} < 0$, we let (5.13b)$\times(-1)$+(5.12a) and (5.12a)$\times(-1)$+(5.12b), then we get

$$\langle k \rangle^{\frac{1}{1-\alpha}} \gtrsim \langle k^{(1)} \rangle^{\frac{y}{1-\alpha}}, \quad \langle k \rangle^{\frac{1}{1-\alpha}} \lesssim \langle k^{(1)} \rangle^{\frac{y}{1-\alpha}},$$

which imply (5.14). Let $\widetilde{k} = (k_1, ..., k_{j-1}, \tilde{k}_j, k_{j+1}, ..., k_n) \in \Lambda(k^{(1)}, k^{(2)})$. In view of (5.5a) and (5.5b) and $(k^{(1)}, k^{(2)}) \in \Omega_{0,2}$, we have

$$\Big| \langle k \rangle^{\frac{\alpha}{1-\alpha}} k_j - \langle \widetilde{k} \rangle^{\frac{\alpha}{1-\alpha}} \widetilde{k}_j \Big| \lesssim \langle k^{(1)} \rangle^{\frac{\alpha}{1-\alpha}} + \langle k \rangle^{\frac{\alpha}{1-\alpha}} + \langle \widetilde{k} \rangle^{\frac{\alpha}{1-\alpha}}.$$

Thus Taylor's theorem, combined with (5.14), gives

$$\Big| k_j - \widetilde{k}_j \Big| \lesssim \langle k^{(1)} \rangle^{\frac{\alpha(1-y)}{1-\alpha}}. \tag{5.15}$$

For $\big(k^{(1)}, k^{(2)}\big) \in \Omega_1 \cup \Omega_2$, in view of (5.6) we have

$$\langle k \rangle \sim \langle k^{(1)} \rangle \vee \langle k^{(2)} \rangle \tag{5.16}$$

and

$$\#\Lambda\big(k^{(1)},k^{(2)}\big)\sim 1. \tag{5.17}$$

In what follows, we separate the proof into four steps. In Steps 1-3, we prove (5.1) for certain p and q. In Step 4, applying the complex interpolation together with the conclusions obtained in the previous three steps, we can get (5.1).

Step 1. $1\leqslant p\leqslant\infty, q\leqslant 1$. Suppose $f,g\in M^{s,\alpha}_{p,q}$, from the triangle inequality and the embedding $\ell^q\subset\ell^1$, we have

$$\begin{aligned}\|fg\|_{M^{s,\alpha}_{p,q}}&=\left(\sum_{k\in\mathbb{Z}^n}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box^\alpha_k(fg)\|^q_p\right)^{\frac{1}{q}}\\&\leqslant\left[\sum_k\langle k\rangle^{\frac{sq}{1-\alpha}}\left(\sum_{k^{(1)},k^{(2)}}\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}}f\Box^\alpha_{k^{(2)}}g)\|_p\right)^q\right]^{\frac{1}{q}}\\&\leqslant\left(\sum_k\langle k\rangle^{\frac{sq}{1-\alpha}}\sum_{k^{(1)},k^{(2)}}\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}}f\Box^\alpha_{k^{(2)}}g)\|^q_p\right)^{\frac{1}{q}}\\&=\left(\sum_{\ell=0}^2\sum_{(k^{(1)},k^{(2)})\in\Omega_\ell}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}}f\Box^\alpha_{k^{(2)}}g)\|^q_p\right)^{\frac{1}{q}}.\end{aligned}\tag{5.18}$$

Applying the multiplier estimate and Hölder's inequality, we see that

$$\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}}f\Box^\alpha_{k^{(2)}}g)\|_p\lesssim\|\Box^\alpha_{k^{(1)}}f\Box^\alpha_{k^{(2)}}g\|_p\lesssim\|\Box^\alpha_{k^{(1)}}f\|_p\|\Box^\alpha_{k^{(2)}}g\|_\infty. \tag{5.19}$$

For $(k^{(1)},k^{(2)})\in\Omega_{0,1}$, when $p=1$ and $s\geqslant n\alpha+\frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{q}-1\big)$, we have

$$\begin{aligned}&\sum_{(k^{(1)},k^{(2)})\in\Omega_{0,1}}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}}f\Box^\alpha_{k^{(2)}}g)\|^q_1\\&\quad\lesssim\sum_{(k^{(1)},k^{(2)})}\langle k^{(1)}\rangle^{\frac{sq\alpha}{1-\alpha}+n\alpha}\|\Box^\alpha_{k^{(1)}}f\|^q_1\langle k^{(2)}\rangle^{\frac{nq\alpha}{1-\alpha}}\|\Box^\alpha_{k^{(2)}}g\|^q_1\leqslant\|f\|^q_{M^{s,\alpha}_{1,q}}\|g\|^q_{M^{s,\alpha}_{1,q}}.\end{aligned}\tag{5.20}$$

If $p=2$ and $s\geqslant\frac{n\alpha}{2}+\frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{q}-1\big)$, by Plancherel and Jensen's inequality*, we have

$$\begin{aligned}&\sum_{(k^{(1)},k^{(2)})\in\Omega_{0,1}}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}}f\Box_{k^{(2)}}g)\|^q_2\\&\quad\lesssim\sum_{(k^{(1)},k^{(2)})}\sup_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}}f\Box^\alpha_{(2)}g)\|^q_2\\&\quad\lesssim\sum_{(k^{(1)},k^{(2)})}\langle k^{(1)}\rangle^{\frac{sq\alpha}{1-\alpha}}[\#\Lambda(k^{(1)},k^{(2)})]^{1-\frac{q}{2}}\|\Box^\alpha_{k^{(1)}}f\Box^\alpha_{k^{(2)}}g\|^q_2\\&\quad\lesssim\sum_{(k^{(1)},k^{(2)})}\langle k^{(1)}\rangle^{\frac{sq\alpha}{1-\alpha}+n\alpha\left(1-\frac{q}{2}\right)}\|\Box^\alpha_{k^{(1)}}f\|^q_2\langle k^{(2)}\rangle^{\frac{nq\alpha}{2(1-\alpha)}}\|\Box^\alpha_{k^{(2)}}g\|^q_2\leqslant\|f\|^q_{M^{s,\alpha}_{2,q}}\|g\|^q_{M^{s,\alpha}_{2,q}}.\end{aligned}\tag{5.21}$$

---

* $a_1^\theta+\ldots+a_N^\theta\leq N^{1-\theta}(a_1+\ldots+a_N)^\theta$ for $\theta\in(0,1)$

For $p=\infty$ and $s \geqslant \frac{n\alpha(1-\alpha)}{2-\alpha}\frac{1}{q}$, we have

$$\begin{aligned}&\sum_{(k^{(1)},k^{(2)})\in\Omega_{0,1}}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g)\|_\infty^q\\&\lesssim\sum_{(k^{(1)},k^{(2)})}\langle k^{(1)}\rangle^{\frac{sq\alpha}{1-\alpha}+n\alpha}\|\Box_{k^{(1)}}^\alpha f\|_\infty^q\|\Box_{k^{(2)}}^\alpha g\|_\infty^q\leqslant\|f\|^q_{M^{s,\alpha}_{\infty,q}}\|g\|^q_{M^{s,\alpha}_{\infty,q}}.\end{aligned}\tag{5.22}$$

For $(k^{(1)},k^{(2)})\in\Omega_{0,2}$, when $s\geqslant n\alpha(1+(1-\alpha)/q)/(2-\alpha)$, we see that

$$2s\geqslant sy+n\alpha\frac{(1+q-y)}{q}.$$

It follows that

$$\begin{aligned}&\sum_{(k^{(1)},k^{(2)})\in\Omega_{0,2}}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g)\|_1^q\\&\lesssim\sum_{k^{(1)},k^{(2)}}\langle k^{(1)}\rangle^{\frac{syq}{1-\alpha}+\frac{n\alpha}{1-\alpha}(1-y)}\|\Box_{k^{(1)}}^\alpha f\|_1^q\langle k^{(2)}\rangle^{\frac{n\alpha q}{1-\alpha}}\|\Box_{k^{(2)}}^\alpha g\|_1^q\lesssim\|f\|_{M^{s,\alpha}_{1,q}}\|g\|_{M^{s,\alpha}_{1,q}}.\end{aligned}\tag{5.23}$$

When $p=\infty$ and $s\geqslant\frac{n\alpha(1-\alpha)}{q(2-\alpha)}$, it is suffices to get

$$\begin{aligned}&\sum_{(k^{(1)},k^{(2)})\in\Omega_{0,2}}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g)\|_\infty^q\\&\lesssim\sum_{k^{(1)},k^{(2)}}\langle k^{(1)}\rangle^{\frac{syq}{1-\alpha}+\frac{n\alpha}{1-\alpha}(1-y)}\|\Box_{k^{(1)}}^\alpha f\|_\infty\|\Box_{k^{(2)}}^\alpha g\|_\infty\lesssim\|f\|_{M^{s,\alpha}_{\infty,q}}\|g\|_{M^{s,\alpha}_{\infty,q}};\end{aligned}\tag{5.24}$$

If $p=2$ and $s\geqslant\frac{n\alpha(\alpha/2+(1-\alpha)/q)}{2-\alpha}$, by Plancherel and Jensen's inequality,

$$\begin{aligned}&\sum_{(k^{(1)},k^{(2)})\in\Omega_{0,2}}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g)\|_2^q\\&\lesssim\sum_{k^{(1)},k^{(2)}}\langle k^{(1)}\rangle^{\frac{syq}{1-\alpha}}[\#\Lambda(k^{(1)},k^{(2)})]^{1-q/2}\|\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g\|_2^q\\&\lesssim\sum_{k^{(1)},k^{(2)}}\langle k^{(1)}\rangle^{\frac{syq}{1-\alpha}+\frac{n\alpha}{(1-\alpha)}(1-y)(1-q/2)}\|\Box_{k^{(1)}}^\alpha f\|_2^q\langle k^{(2)}\rangle^{\frac{n\alpha q}{2(1-\alpha)}}\|\Box_{k^{(2)}}^\alpha g\|_2^q\\&\lesssim\|f\|_{M^{s,\alpha}_{2,q}}\|g\|_{M^{s,\alpha}_{2,q}}.\end{aligned}\tag{5.25}$$

From (5.16), (5.17), (2.2), if $(k^{(1)},k^{(2)})\in\Omega_1$, when $s\geqslant\frac{n\alpha}{p}$,

$$\begin{aligned}&\sum_{(k^{(1)},k^{(2)})\in\Omega_1}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g)\|_p^q\\&\lesssim\sum_{k^{(1)}\in\mathbb{Z}^n}\langle k^{(1)}\rangle^{\frac{sq}{1-\alpha}}\|\Box_{k^{(1)}}^\alpha f\|_p^q\sum_{k^{(2)}\in\mathbb{Z}^n}\|\Box_{k^{(2)}}^\alpha g\|_\infty^q\lesssim\|f\|_{M^{s,\alpha}_{p,q}}\|g\|_{M^{0,\alpha}_{\infty,q}}\\&\lesssim\|f\|_{M^{s,\alpha}_{p,q}}\|g\|_{M^{s,\alpha}_{p,q}}.\end{aligned}\tag{5.26}$$

Similarly, if $(k^{(1)},k^{(2)})\in\Omega_2$ and $s\geqslant\frac{n\alpha}{p}$,

$$\sum_{(k^{(1)},k^{(2)})\in\Omega_2}\sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g)\|_p^q\lesssim\|f\|_{M^{s,\alpha}_{p,q}}\|g\|_{M^{s,\alpha}_{p,q}}.\tag{5.27}$$

By complex interpolation, (5.18)-(5.27) imply that $M_{p,q}^{s,\alpha}$ is a multiplication algebra as long as $s \geqslant s_0$ for some s_0. More precisely, when $1 \leqslant p \leqslant 2$, from (5.20) and (5.21), we get $s_0 = \frac{n\alpha}{p} + \frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{q} - 1\big)$; and when $2 < p \leqslant \infty$, from (5.21) and (5.22), we get $s_0 = \frac{n\alpha}{p} + \frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{q} - \frac{2}{p}\big)$.

Step 2. $0 < p \leqslant \infty, q = \infty$. First, we consider the case $1 \leqslant p \leqslant \infty$. Suppose $f, g \in M_{p,\infty}^{s,\alpha}$, from the triangle inequality, we have

$$\begin{aligned}
\|fg\|_{M_{p,\infty}^{s,\alpha}} &= \sup_{k\in\mathbb{Z}^n} \langle k\rangle^{\frac{s}{1-\alpha}} \|\Box_k^\alpha(fg)\|_p \\
&\leqslant \sup_{k\in\mathbb{Z}^n} \langle k\rangle^{\frac{s}{1-\alpha}} \sum_{(k^{(1)},k^{(2)})\in\Lambda(k)} \|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f \Box_{k^{(2)}}^\alpha g)\|_p \\
&= \sup_{k\in\mathbb{Z}^n} \sum_{\ell=0}^{2} \sum_{(k^{(1)},k^{(2)})\in\Lambda(k)\cap\Omega_\ell} \langle k\rangle^{\frac{s}{1-\alpha}} \|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f \Box_{k^{(2)}}^\alpha g)\|_p.
\end{aligned} \tag{5.28}$$

For a $\Psi \subset \mathbb{Z}^{2n}$, we denote

$$\begin{aligned}
\Psi_1^\perp &= \Big\{k^{(1)} \in \mathbb{Z}^n : \exists k^{(2)} \in \mathbb{Z}^n \text{ s.t. } \big(k^{(1)}, k^{(2)}\big) \in \Psi\Big\}; \\
\Psi_2^\perp &= \Big\{k^{(2)} \in \mathbb{Z}^n : \exists k^{(1)} \in \mathbb{Z}^n \text{ s.t. } \big(k^{(1)}, k^{(2)}\big) \in \Psi\Big\}.
\end{aligned}$$

Let $s > \frac{n\alpha}{p} + n(1-\alpha)$. For any $k^{(2)} \in \{\{\Omega_0 \cup \Omega_1\} \cap \Lambda(k)\}_2^\perp$ with every fixed k, noticing (5.6), we easily see $\#\Lambda(-k^{(2)}, k) \lesssim 1$. Inserting (5.19) and using (2.3), we obtain

$$\begin{aligned}
&\sum_{(k^{(1)},k^{(2)})\in\{\Omega_0\cup\Omega_1\}\cap\Lambda(k)} \langle k\rangle^{\frac{s}{1-\alpha}} \|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f \Box_{k^{(2)}}^\alpha g)\|_p \\
&\lesssim \sup_{k^{(1)}\in\big\{\{\Omega_0\cup\Omega_1\}\cap\Lambda(k)\big\}_1^\perp} \langle k\rangle^{\frac{s}{1-\alpha}} \|\Box_{k^{(1)}}^\alpha f\|_p \sum_{k^{(1)}\in\big\{\{\Omega_0\cup\Omega_1\}\cap\Lambda(k)\big\}_1^\perp} \sum_{k^{(2)}\in\Lambda(-k^{(1)},k)} \|\Box_{k^{(2)}}^\alpha g\|_\infty \\
&\lesssim \sup_{k^{(1)}\in\mathbb{Z}^n} \langle k^{(1)}\rangle^{\frac{s}{1-\alpha}} \|\Box_{k^{(1)}}^\alpha f\|_p \sum_{k^{(2)}\in\big\{\{\Omega_0\cup\Omega_1\}\cap\Lambda(k)\big\}_2^\perp} \sum_{k^{(1)}\in\Lambda(-k^{(2)},k)} \|\Box_{k^{(2)}}^\alpha g\|_\infty \\
&\lesssim \|f\|_{M_{p,\infty}^{s,\alpha}} \|g\|_{M_{\infty,1}^{0,\alpha}} \lesssim \|f\|_{M_{p,\infty}^{s,\alpha}} \|g\|_{M_{p,\infty}^{s,\alpha}}.
\end{aligned} \tag{5.29}$$

For $k^{(1)} \in \{\Omega_2 \cap \Lambda(k)\}_1^\perp$ with every fixed k, symmetrically, we have

$$\begin{aligned}
&\sum_{(k^{(1)},k^{(2)})\in\Omega_2\cap\Lambda(k)} \langle k\rangle^{\frac{s}{1-\alpha}} \|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f \Box_{k^{(2)}}^\alpha g)\|_p \\
&\quad\lesssim \sup_{k^{(2)}\in\mathbb{Z}^n} \langle k^{(2)}\rangle^{\frac{s}{1-\alpha}} \|\Box_{k^{(2)}}^\alpha g\|_p \sum_{k^{(1)}\in\{\Omega_2\cap\Lambda(k)\}_1^\perp} \sum_{k^{(2)}\in\Lambda(-k^{(1)},k)} \|\Box_{k^{(1)}}^\alpha f\|_\infty \\
&\quad\lesssim \|f\|_{M_{\infty,1}^{0,\alpha}} \|g\|_{M_{p,\infty}^{s,\alpha}} \lesssim \|f\|_{M_{p,\infty}^{s,\alpha}} \|g\|_{M_{p,\infty}^{s,\alpha}}.
\end{aligned} \tag{5.30}$$

Combining (5.28)–(5.30), we know when $s > \frac{n\alpha}{p} + n(1-\alpha)$, $M_{p,\infty}^{s,\alpha}$ is a multiplication algebra.

Next, we consider the case $0 < p < 1$ and $q = \infty$. Suppose that $f, g \in M_{p,\infty}^{s,\alpha}$. It follows from the embedding $\ell^p \subset \ell^1$ that

$$\begin{aligned}\|fg\|_{M^{s,\alpha}_{p,\infty}} &\leqslant \sup_k \langle k\rangle^{\frac{s}{1-\alpha}} \left(\sum_{(k^{(1)},k^{(2)})\in\Lambda(k)} \|\Box^\alpha_k(\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g)\|_p^p\right)^{\frac{1}{p}} \\ &= \sup_k \left(\sum_{\ell=0}^{2} \sum_{(k^{(1)},k^{(2)})\in\Lambda(k)\cap\Omega_\ell} \langle k\rangle^{\frac{sp}{1-\alpha}} \|\Box^\alpha_k(\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g)\|_p^p\right)^{\frac{1}{p}}.\end{aligned} \tag{5.31}$$

By Proposition 2.1, for $(k^{(1)},k^{(2)}) \in \Lambda(k)\cap\Omega_0$, we have

$$\begin{aligned}\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g)\|_p &\lesssim \langle k^{(1)}\rangle^{\frac{n\alpha}{(1-\alpha)}(\frac{1}{p}-1)} \|\mathscr{F}^{-1}\eta^\alpha_k\|_p \|\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g\|_p \\ &\lesssim \langle k^{(1)}\rangle^{\frac{n\alpha}{(1-\alpha)}(\frac{1}{p}-1)} \langle k\rangle^{-\frac{n\alpha}{(1-\alpha)}(\frac{1}{p}-1)} \|\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g\|_p.\end{aligned} \tag{5.32}$$

When $s \geq \frac{n}{p} + \frac{n\alpha(1-\alpha)}{(2-\alpha)}(\frac{1}{p}-1)$, inserting (5.32) and using (2.3), we obtain that

$$\begin{aligned}&\sum_{(k^{(1)},k^{(2)})\in\Lambda(k)\cap\Omega_0} \langle k\rangle^{\frac{sp}{1-\alpha}} \|\Box^\alpha_k(\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g)\|_p^p \\ &\lesssim \sum_{(k^{(1)},k^{(2)})\in\Lambda(k)\cap\Omega_0} \langle k^{(1)}\rangle^{\frac{spy+n\alpha(1-y)(1-p)}{1-\alpha}} \|\Box^\alpha_{k^{(1)}} f\|_p^p \|\Box^\alpha_{k^{(2)}} g\|_\infty^p \\ &\lesssim \sup_{k^{(1)}\in\mathbb{Z}^n} \langle k^{(1)}\rangle^{\frac{sp}{1-\alpha}} \|\Box^\alpha_{k^{(1)}} f\|_p^p \sum_{k^{(2)}\in\{\Lambda(k)\cap\Omega_0\}^{\perp}_{2}} \sum_{k^{(1)}\in\Lambda(-k^{(2)},k)} \langle k^{(2)}\rangle^{\frac{sp(y-1)+n\alpha(1-y)(1-p)}{1-\alpha}} \|\Box^\alpha_{k^{(2)}} g\|_\infty^p \\ &\lesssim \|f\|^p_{M^{s,\alpha}_{p,\infty}} \|g\|^p_{M^{s,\alpha}_{p,\infty}}.\end{aligned} \tag{5.33}$$

For any $(k^{(1)},k^{(2)}) \in \Lambda(k)\cap\{\Omega_1\cup\Omega_2\}$ with every fixed k, imitating the process as in (5.32) and combining (5.16), we get

$$\|\Box^\alpha_k(\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g)\|_p \lesssim \|\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g\|_p. \tag{5.34}$$

When $s > \frac{n}{p}$, inserting (5.34) and also using (2.3), we obtain

$$\begin{aligned}&\sum_{\ell=1}^{2} \sum_{(k^{(1)},k^{(2)})\in\Lambda(k)\cap\Omega_\ell} \langle k\rangle^{\frac{sp}{1-\alpha}} \|\Box^\alpha_k(\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g)\|_p^p \\ &\lesssim \|f\|^p_{M^{s,\alpha}_{p,\infty}} \|g\|^p_{M^{0,\alpha}_{\infty,p}} + \|f\|^p_{M^{0,\alpha}_{\infty,p}} \|g\|^p_{M^{s,\alpha}_{p,\infty}} \lesssim \|f\|^p_{M^{s,\alpha}_{p,\infty}} \|g\|^p_{M^{s,\alpha}_{p,\infty}}.\end{aligned} \tag{5.35}$$

Combining (5.31),(5.33) and (5.35), we conclude when $s \geq \frac{n}{p} + \frac{n\alpha(1-\alpha)}{(2-\alpha)}((1\vee\frac{1}{p})-1)$, $M^{s,\alpha}_{p,\infty}$ is a multiplication algebra.

Step 3. $p < 1,\ q = p$. Suppose $f, g \in M^{s,\alpha}_{p,p}$. From the embedding $\ell^p \subset \ell^1 \subset \ell^{1/p}$ it follows that

$$\begin{aligned}\|fg\|_{M^{s,\alpha}_{p,p}} &\leqslant \left(\sum_k \langle k\rangle^{\frac{sp}{1-\alpha}} \sum_{k^{(1)},k^{(2)}} \|\Box^\alpha_k(\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g)\|_p^p\right)^{\frac{1}{p}} \\ &= \left(\sum_{\ell=0}^{2} \sum_{(k^{(1)},k^{(2)})\in\Omega_\ell} \sum_{k\in\Lambda(k^{(1)},k^{(2)})} \langle k\rangle^{\frac{sp}{1-\alpha}} \|\Box^\alpha_k(\Box^\alpha_{k^{(1)}} f \Box^\alpha_{k^{(2)}} g)\|_p^p\right)^{\frac{1}{p}}.\end{aligned} \tag{5.36}$$



Figure 5: For Step 4 in the proof of Theorem 5.1.

For $(k^{(1)},k^{(2)})\in\Omega_0$, if $s\geqslant\frac{n\alpha}{p}+\frac{n\alpha(1-\alpha)}{(2-\alpha)}(\frac{1}{p}-1)$, then we see from (5.10), (5.14), (5.15) and (5.32) that

$$
\begin{aligned}
&\sum_{(k^{(1)},k^{(2)})\in\Omega_0}\ \sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sp}{1-\alpha}}\|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g)\|_p^p\\
&\quad\lesssim\sum_{(k^{(1)},k^{(2)})\in\Omega_0}\langle k^{(1)}\rangle^{\frac{spy}{1-\alpha}-\frac{n\alpha y}{1-\alpha}(1-p)+\frac{n\alpha}{1-\alpha}(1-y)+\frac{n\alpha}{1-\alpha}(1-p)}\|\Box_{k^{(1)}}^\alpha f\|_p^p\|\Box_{k^{(2)}}^\alpha g\|_\infty^p\\
&\quad\lesssim\sum_{(k^{(1)},k^{(2)})\in\Omega_0}\langle k^{(1)}\rangle^{\frac{spy}{1-\alpha}+\frac{n\alpha}{1-\alpha}(1-p)(1-y)+\frac{n\alpha}{1-\alpha}(2-y)}\|\Box_{k^{(1)}}^\alpha f\|_p^p\|\Box_{k^{(2)}}^\alpha g\|_p^p\\
&\quad\lesssim\|f\|_{M_{p,p}^{s,\alpha}}^p\|g\|_{M_{p,p}^{s,\alpha}}^p.
\end{aligned}
\tag{5.37}
$$

For $(k^{(1)},k^{(2)})\in\Omega_1\cup\Omega_2$, when $s\geqslant\frac{n\alpha}{p}$, in view of (2.2), we obtain

$$
\begin{aligned}
&\sum_{(k^{(1)},k^{(2)})\in\Omega_1\cup\Omega}\ \sum_{k\in\Lambda(k^{(1)},k^{(2)})}\langle k\rangle^{\frac{sp}{1-\alpha}}\|\Box_k^\alpha(\Box_{k^{(1)}}^\alpha f\Box_{k^{(2)}}^\alpha g)\|_p^p\\
&\quad\lesssim\|f\|_{M_{p,p}^{s,\alpha}}^p\|g\|_{M_{\infty,p}^{0,\alpha}}^p+\|f\|_{M_{\infty,p}^{0,\alpha}}^p\|g\|_{M_{p,p}^{s,\alpha}}^p\lesssim\|f\|_{M_{p,p}^{s,\alpha}}^p\|g\|_{M_{p,p}^{s,\alpha}}^p.
\end{aligned}
\tag{5.38}
$$

Combining (5.36)-(5.38), we conclude when $s\geqslant\frac{n\alpha}{p}+\frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{p}-1\big)$, $M_{p,p}^{s,\alpha}$ is a multiplication algebra.

Step 4. Let $\big(\frac{1}{p},\frac{1}{q}\big)\in\big\{\big(\frac{1}{p},\frac{1}{q}\big)\in \mathrm{D}_1:\frac{1}{q}<1\big\}$. It is easy to see that $\big(\frac{1}{p},\frac{1}{q}\big)$ is a point at the line segment connecting $\big(\frac{1}{p}+\frac{1}{2}\big(1-\frac{1}{q}\big),1\big)$ and $\big(0,\frac{1}{q}-\frac{2}{p}\big)$, which is parallel to the line connecting $\big(\frac{1}{2},1\big)$ and $(0,0)$. At the point $(\frac{1}{\bar{p}},1):=\big(\frac{1}{p}+\frac{1}{2}\big(1-\frac{1}{q}\big),1\big)$, in Step 1 we have shown that $M_{\bar{p},1}^{s,\alpha}$ is a multiplication algebra if $s\geqslant n\alpha\big[\frac{1}{p}+\frac{1}{2}\big(1-\frac{1}{q}\big)\big]+\frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{q}-\frac{2}{p}\big)$. For $(0,\frac{1}{\bar{q}}):=\big(0,\frac{1}{q}-\frac{2}{p}\big)$, complex interpolation between $(0,1)$ in Step 1 and $(0,0)$ in Step 3 shows that once $s > n(1-\alpha)\big(1-\frac{1}{q}+\frac{2}{p}\big)+\frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{q}-\frac{2}{p}\big)$, the associated α-modulation space $M^{s,\alpha}_{\infty,\bar{q}}$ is a multiplication algebra. Again, using the complex interpolation and combining the result in Step 2, and we arrive at (5.1) in D_1. Denote (see Fig. 5)

$$\mathrm{D}_{21} = [0,1]\times[0,1]\backslash \mathrm{D}_1;$$
$$\mathrm{D}_{23} = \Big\{\big(\tfrac{1}{p},\tfrac{1}{q}\big)\in\mathbb{R}^2_+ : \tfrac{1}{q}\geqslant\tfrac{1}{p}\Big\}\Big\backslash \mathrm{D}_1\cup \mathrm{D}_{21}.$$

Through the point $\big(\frac{1}{p},\frac{1}{q}\big)\in \mathrm{D}_{21}$, one can make a line segment connecting $\big(\frac{1}{2q},\frac{1}{q}\big)$ and $\big(1,\frac{1}{q}\big)$. For $\big(\frac{1}{2q},\frac{1}{q}\big)$, we see that once $s>\frac{n\alpha}{2q}+n(1-\alpha)\big(1-\frac{1}{q}\big)$, $M^{s,\alpha}_{2q,q}$ is a multiplication algebra. For $\big(1,\frac{1}{q}\big)$, complex interpolation between $(1,1)$ in Step 1 and $(1,0)$ in Step 3 shows that once $s>n\alpha+n(1-\alpha)\big(1-\frac{1}{q}\big)$, the associated α-modulation space is a multiplication algebra. Then we use complex interpolation to get that $M^{s,\alpha}_{p,q}$ is a multiplication algebra if $s>\frac{n\alpha}{p}+n(1-\alpha)\big(1-\frac{1}{q}\big)$. If $(1/p,1/q)\in \mathrm{D}_{23}$, the result can be derived in a similar way.

If $\big(\frac{1}{p},\frac{1}{q}\big)\in \mathrm{D}_{22}$, then it belongs to the segment by connecting $\big(\frac{1}{p},0\big)$ and $\big(\frac{1}{p},\frac{1}{p}\big)$. In Step 4 we see that once $s\geqslant\frac{n\alpha}{p}+n(1-\alpha)\frac{1}{p}+\frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{p}-1\big)$, $M^{s,\alpha}_{p,\infty}$ is a multiplication algebra; and once $s\geqslant\frac{n\alpha}{p}+\frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{p}-1\big)$, $M^{s,\alpha}_{p,p}$ is a multiplication algebra. Then complex interpolation between them gives once $s\geqslant\frac{n\alpha}{p}+n(1-\alpha)\big(\frac{1}{p}-\frac{1}{q}\big)+\frac{n\alpha(1-\alpha)}{2-\alpha}\big(\frac{1}{p}-1\big)$, $M^{s,\alpha}_{p,q}$ is a multiplication algebra. $\square$

6 Sharpness for the scaling and embedding properties

In this section we show the necessity of Theorems 3.1, 4.1 and 4.2. Since the *p-BAPU* has no scaling, it is difficult to calculate the norm for a known function. However, we have the following equivalent norm on α-modulation spaces. Let ρ be a smooth radial bump function supported in $B(0,2)$, satisfying $\rho(\xi)=1$ as $|\xi|<1$, and $\rho(\xi)=0$ as $|\xi|\geqslant 2$. Let ρ^α_k be as in (1.4):

$$\rho^\alpha_k(\xi)=\rho\left(\frac{\xi-\langle k\rangle^{\alpha/(1-\alpha)}k}{C\langle k\rangle^{\alpha/(1-\alpha)}}\right).$$

Proposition 6.1. *Let ρ^α_k be as in the above. Then*

$$\|f\|^{\circ}_{M^{s,\alpha}_{p,q}}=\left(\sum_{k\in\mathbb{Z}^n}\langle k\rangle^{\frac{sq}{1-\alpha}}\|(\mathscr{F}^{-1}\rho^\alpha_k)*f\|^q_p\right)^{1/q}$$

is an equivalent norm on $M^{s,\alpha}_{p,q}$.

Proof. If $p\geq 1$, in view of Young's inequality,

$$\|(\mathscr{F}^{-1}\rho^\alpha_k)*f\|_p\leq\sum_{|\ell|_\infty\leq C}\|(\mathscr{F}^{-1}\rho^\alpha_{k+\ell})\|_1\|\Box^\alpha_k f\|_p\lesssim\|\Box^\alpha_k f\|_p.$$

If $p<1$, by Proposition 2.1 and the scaling argument, we have

$$\|(\mathscr{F}^{-1}\rho^\alpha_k)*f\|_p\leq\langle k\rangle^{n\alpha(1/p-1)/(1-\alpha)}\sum_{|\ell|_\infty\leq C}\|(\mathscr{F}^{-1}\rho^\alpha_{k+\ell})\|_p\|\Box^\alpha_k f\|_p\lesssim\|\Box^\alpha_k f\|_p.$$

Combining the above two cases, we have $\|f\|^\circ_{M^{s,\alpha}_{p,q}} \le \|f\|_{M^{s,\alpha}_{p,q}}$. On the other hand, noticing that

$$\eta^\alpha_k = \rho^\alpha_k \frac{\rho\left(\frac{\xi-\langle k\rangle^{\alpha/(1-\alpha)}k}{2C\langle k\rangle^{\alpha/(1-\alpha)}}\right)}{\sum_{\ell\in\mathbb{Z}^n}\rho\left(\frac{\xi-\langle k+\ell\rangle^{\alpha/(1-\alpha)}(k+\ell)}{C\langle k+\ell\rangle^{\alpha/(1-\alpha)}}\right)} := \rho^\alpha_k\sigma^\alpha_k,$$

We have for $p \ge 1$, in view of Young's inequality,

$$\|\Box^\alpha_k f\|_p \le \|(\mathscr{F}^{-1}\rho^\alpha_k) * f\|_p \|(\mathscr{F}^{-1}\sigma^\alpha_k)\|_1 \lesssim \|(\mathscr{F}^{-1}\rho^\alpha_k) * f\|_p.$$

If $p < 1$, by Proposition 2.1, the scaling argument and the generalized Bernstein inequality, we have

$$\|\Box^\alpha_k f\|_p \lesssim \langle k\rangle^{n\alpha(1/p-1)/(1-\alpha)} \|(\mathscr{F}^{-1}\rho^\alpha_k) * f\|_p \|\mathscr{F}^{-1}\sigma^\alpha_k\|_p \|\Box^\alpha_k f\|_p \lesssim \|(\mathscr{F}^{-1}\rho^\alpha_k) * f\|_p.$$

The result follows. $\Box$

Proof of Theorem 3.1. (**Necessity**) We divide the proof into the following two cases $\lambda \gg 1$ and $\lambda \ll 1$, respectively.

Case 1. $\lambda \gg 1$. One needs to show that

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \gtrsim \lambda^{-\frac{n}{p}+n\left(\frac{1}{p}-1\right)\vee(s+s_c)} \|f\|_{M^{s,\alpha}_{p,q}}.$$

Case 1.1. We consider the case $s_p = n(1/p-1) > s + s_c$. Our aim is to show that there exists a function f satisfying

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \gtrsim \lambda^{-n}\|f\|_{M^{s,\alpha}_{p,q}}.$$

Taking $f = \mathscr{F}^{-1}\rho$, we have

$$\|f_\lambda\|^\circ_{M^{s,\alpha}_{p,q}} \ge \|(\mathscr{F}^{-1}\rho^\alpha_0) * f_\lambda\|_p \geqslant \lambda^{-n}\|\mathscr{F}^{-1}\rho\|_p \gtrsim \lambda^{-n}\|f\|_{M^{s,\alpha}_{p,q}}. \tag{6.1}$$

Case 1.2. We consider the case $s_p < s + s_c$. According to the definition of s_c, we separate the proof into the following three cases.

Case 1.2.1. $s_c = n(1-\alpha)(1/p + 1/q - 1)$. Put $f = \mathscr{F}^{-1}\rho$. Since $\lambda \gg 1$, we see that for some $0 < \varepsilon_0 < \varepsilon_1 \ll 1$,

$$(\mathscr{F}^{-1}\rho^\alpha_k) * f_\lambda = \lambda^{-n}\mathscr{F}^{-1}\rho^\alpha_k, \qquad |k| \in [\varepsilon_0\lambda^{1-\alpha},\ \varepsilon_1\lambda^{1-\alpha}].$$

It follows that

$$\begin{aligned}\|f_\lambda\|^\circ_{M^{s,\alpha}_{p,q}} &\gtrsim \lambda^{-n}\left(\sum_{|l|\in[\varepsilon_0\lambda^{1-\alpha},\ \varepsilon_1\lambda^{1-\alpha}]} \langle l\rangle^{\frac{sq}{1-\alpha}}\|\mathscr{F}^{-1}\rho^\alpha_l\|^q_p\right)^{\frac{1}{q}} \\ &\gtrsim \lambda^{-\frac{n}{p}+s+n(1-\alpha)\left(\frac{1}{p}+\frac{1}{q}-1\right)} \\ &\gtrsim \lambda^{-\frac{n}{p}+s+n(1-\alpha)\left(\frac{1}{p}+\frac{1}{q}-1\right)}\|f\|_{M^{s,\alpha}_{p,q}}.\end{aligned} \tag{6.2}$$

Case 1.2.2. $s_c = 0$. Let us take $f = \mathrm{e}^{\mathrm{i}x_1}\mathscr{F}^{-1}\rho_\lambda$. We have $f_\lambda = \lambda^{-n}e^{\mathrm{i}x_1\lambda}\mathscr{F}^{-1}\rho$. We may assume that there exists $l_0 \in \mathbb{N}$ such that $\langle l_0\rangle^{\alpha/(1-\alpha)}l_0 \sim \lambda$ and $B((\lambda,0,...,0),\ 2) \subset B(\langle l_0\rangle^{\alpha/(1-\alpha)}(l_0,0,...,0),\ C\langle l_0\rangle^{\alpha/(1-\alpha)})$. It is easy to see that $\|f\|_p \sim \lambda^{n\left(\frac{1}{p}-1\right)}$ and

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \sim \lambda^{-n}\langle l_0\rangle^{s/(1-\alpha)} \sim \lambda^{-n+s} \gtrsim \lambda^{-\frac{n}{p}+s}\|f\|_p \gtrsim \lambda^{-\frac{n}{p}+s}\|f\|_{M^{s,\alpha}_{p,q}}. \tag{6.3}$$

Case 1.2.3. $s_c = n(1-\alpha)(1/q - 1/p)$. Put

$$f^{(l)} = \mathrm{e}^{\mathrm{i}x\langle l\rangle^{\frac{\alpha}{1-\alpha}} l/\lambda}\ \tau_{\lambda^{1-\alpha}l}\left(\rho\left(\frac{\lambda\ \cdot}{c\langle l\rangle^{\frac{\alpha}{1-\alpha}}}\right)\right)^{\vee}, \quad f = \sum_{|\ell|\sim\lambda^{1-\alpha}} f^{(l)}, \tag{6.4}$$

where $|\ell| \sim \lambda^{1-\alpha}$ means that $|l| \in [\varepsilon_0\lambda^{1-\alpha},\ \varepsilon_1\lambda^{1-\alpha}]$ for some $0 < \varepsilon_0 < \varepsilon_1 \ll 1$. If $|l| \sim \lambda^{1-\alpha}$, then $\|f^{(l)}\|_p \sim \lambda^{n(1-\alpha)\left(\frac{1}{p}-1\right)}$, and

$$(f^{(l)})_\lambda = \mathrm{e}^{\mathrm{i}x\langle l\rangle^{\frac{\alpha}{1-\alpha}} l}\ \tau_{\lambda^{1-\alpha}l}\left(\rho\left(\frac{\cdot}{c\langle l\rangle^{\frac{\alpha}{1-\alpha}}}\right)\right)^{\vee}, \tag{6.5}$$

which follows that $(\mathscr{F}^{-1}\rho_l^\alpha) * (f^{(l)})_\lambda = (f^{(l)})_\lambda$. Since supp ρ_l^α overlaps at most finite many supp ρ_{l+k}^α, we see that $\Box_{l+k}^\alpha (f^{(l)})_\lambda = 0$ if $|k| \geq C$. Let $A(\lambda) \subset \{l : |l| \sim \lambda^{1-\alpha}\}$ be the set so that for any $l, \tilde{l} \in A(\lambda)$ $(l \neq \tilde{l})$, $|l - \tilde{l}| \geq C$. We have

$$\begin{aligned}
\|f_\lambda\|_{M^{s,\alpha}_{p,q}} &\gtrsim \left(\sum_{l\in A(\lambda)} \langle l\rangle^{\frac{sq}{1-\alpha}} \|(f^{(l)})_\lambda\|_p^q\right)^{\frac{1}{q}} \\
&\gtrsim \lambda^{-\frac{n}{p}+s}\left(\sum_{l\in A(\lambda)} \|f^{(l)}\|_p^q\right)^{\frac{1}{q}} \\
&\gtrsim \lambda^{-\frac{n}{p}+s+n(1-\alpha)\left(\frac{1}{q}+\frac{1}{p}-1\right)}.
\end{aligned} \tag{6.6}$$

Moreover, we easily see that

$$|\widehat{f^{(l)}}(\xi)| = \rho\left(\frac{\lambda\xi - \langle l\rangle^{\alpha/(1-\alpha)} l}{c\langle l\rangle^{\alpha/(1-\alpha)}}\right).$$

It follows that supp $\widehat{f^{(l)}}$ is included in the unit ball. Hence, we have

$$\|f\|_{M^{s,\alpha}_{p,q}} \lesssim \left\|\sum_{l\in A(\lambda)} f^{(l)}\right\|_p.$$

By Plancheral's identity,

$$\|f\|_2 = \left\|\widehat{f}\right\|_2 = \left(\sum_{l\in A(\lambda)} \int \rho^2\left(\frac{\lambda\xi - \langle l\rangle^{\alpha/(1-\alpha)} l}{c\langle l\rangle^{\alpha/(1-\alpha)}}\right) d\xi\right)^{1/2} \sim 1.$$

On the other hand, in view of $\mathscr{F}^{-1}\rho$ is a Schwartz function, we have

$$|\mathscr{F}^{-1}\rho(x)| \lesssim \langle x\rangle^{-N}.$$

It follows that for $N \gg n$,

$$\left|\sum_{l\in A(\lambda)} f^{(l)}(x)\right| \lesssim \lambda^{-n(1-\alpha)} \sum_{l\in A(\lambda)} \left(1 + \lambda^{-(1-\alpha)}|x - \lambda^{1-\alpha} l|\right)^{-N} \lesssim \lambda^{-n(1-\alpha)}.$$

By Hölder's inequality,

$$\|f\|_{M^{s,\alpha}_{p,q}} \lesssim \|f\|_p \lesssim \lambda^{n(1-\alpha)(2/p-1)}.$$

The result follows.

Case 2. $\lambda \ll 1$. It suffices to show that for some $f \in M^{s,\alpha}_{p,q}$,

$$\|f_\lambda\|^\circ_{M^{s,\alpha}_{p,q}} \gtrsim \lambda^{-\frac{n}{p}}(1 \vee \lambda^{s+s_c})\|f\|_{M^{s,\alpha}_{p,q}}.$$

Case 2.1. $s+s_c \geq 0$. Taking $f = \mathscr{F}^{-1}\rho$, we have

$$\|f_\lambda\|^\circ_{M^{s,\alpha}_{p,q}} = \|(\mathscr{F}^{-1}\rho)(\lambda\,\cdot)\|_p \sim \lambda^{-\frac{n}{p}}\|f\|_{M^{s,\alpha}_{p,q}}. \tag{6.7}$$

Case 2.2. $s+s_c < 0$. We divide the proof into the following three cases.

Case 2.2.1. We can find some k_0 such that $\lambda\langle k_0\rangle^{\frac{1}{1-\alpha}} \sim 1$. Denote

$$f = \mathrm{e}^{\mathrm{i}x\langle k_0\rangle^{\frac{\alpha}{1-\alpha}}k_0}\mathscr{F}^{-1}\left(\rho\left(\frac{\cdot}{c\langle k_0\rangle^{\frac{\alpha}{1-\alpha}}}\right)\right), \tag{6.8}$$

We have

$$\widehat{f_\lambda} = \lambda^{-n}\rho\left(\frac{\xi/\lambda - \langle k_0\rangle^{\frac{\alpha}{1-\alpha}}k_0}{c\langle k_0\rangle^{\frac{\alpha}{1-\alpha}}}\right), \tag{6.9}$$

Therefore,

$$\|f_\lambda\|^\circ_{M^{s,\alpha}_{p,q}} \gtrsim \|f_\lambda\|_p \gtrsim \lambda^{-\frac{n}{p}+s}\langle k_0\rangle^{\frac{s}{1-\alpha}}\|f\|_p \gtrsim \lambda^{-\frac{n}{p}+s}\|f\|_{M^{s,\alpha}_{p,q}}. \tag{6.10}$$

Case 2.2.2. Taking $f = \lambda^n\mathscr{F}^{-1}\rho_\lambda$, we have $f_\lambda = \mathscr{F}^{-1}\rho$. It follows that $\|f_\lambda\|^\circ_{M^{s,\alpha}_{p,q}} \sim 1$. On the other hand,

$$\begin{aligned}
\|f\|^\circ_{M^{s,\alpha}_{p,q}} &\lesssim \left(\sum_{|k|\lesssim\lambda^{\alpha-1}}\langle k\rangle^{\frac{sq}{1-\alpha}}\|(\mathscr{F}^{-1}\rho^\alpha_k)*f\|^q_p\right)^{\frac{1}{q}} \\
&\lesssim \left(\sum_{|k|\lesssim\lambda^{\alpha-1}}\langle k\rangle^{\frac{sq}{1-\alpha}}\|\mathscr{F}^{-1}\rho^\alpha_k)\|^q_p\|f\|^q_1\right)^{\frac{1}{q}} \\
&\lesssim \lambda^{n-s-n(1-\alpha)\frac{1}{q}-n\alpha\left(1-\frac{1}{p}\right)}.
\end{aligned} \tag{6.11}$$

It follows that

$$\|f_\lambda\|^\circ_{M^{s,\alpha}_{p,q}} \gtrsim \lambda^{-\frac{n}{p}+s+s_c}\|f\|_{M^{s,\alpha}_{p,q}}.$$

Case 2.2.3. $s_c = n(1-\alpha)(1/q-1/p)$. Let $A(\lambda)$ be the set so that for any $l,\tilde{l} \in A(\lambda)$ $(l \neq \tilde{l})$, $|l-\tilde{l}| \geq C$ and $|l| \in [\varepsilon_0\lambda^{\alpha-1},\ \varepsilon_1\lambda^{\alpha-1}]$ for some $0 < \varepsilon_0 < \varepsilon_1 \ll 1$. Take

$$f^{(l)} = \mathrm{e}^{\mathrm{i}x\langle l\rangle^{\frac{\alpha}{1-\alpha}}l}\tau_{C'l}\mathscr{F}^{-1}\left(\rho\left(\frac{\cdot}{c\langle l\rangle^{\frac{\alpha}{1-\alpha}}}\right)\right), \quad f = \sum_{l\in A(\lambda)} f^{(l)}. \tag{6.12}$$

One easily sees that

$$(f^{(l)})_\lambda = \lambda^{-n}\mathrm{e}^{\mathrm{i}x\lambda\langle l\rangle^{\frac{\alpha}{1-\alpha}}l}\tau_{C'l}\mathscr{F}^{-1}\left(\rho\left(\frac{\cdot}{c\lambda\langle l\rangle^{\frac{\alpha}{1-\alpha}}}\right)\right). \tag{6.13}$$

We have

$$\begin{aligned}\|f\|^\circ_{M^{s,\alpha}_{p,q}} &= \left(\sum_{l\in A(\lambda)} \langle l\rangle^{\frac{sq}{1-\alpha}} \|(f^{(l)})\|_p^q\right)^{\frac1q} \\ &\lesssim \lambda^{-s}\left(\sum_{l\in A(\lambda)} \langle l\rangle^{\frac{n\alpha}{1-\alpha}(1-\frac1p)q}\right)^{\frac1q} \\ &\lesssim \lambda^{-s+n\alpha\left(\frac1p-1\right)+\frac{n(\alpha-1)}{q}}.\end{aligned} \tag{6.14}$$

Since $\mathrm{supp}\widehat{f}^{(l)}_\lambda$ is contained in the unit ball, we see that

$$\|f_\lambda\|_{M^{s,\alpha}_{p,q}} \gtrsim \left\|\sum_{l\in A(\lambda)} f^{(l)}_\lambda\right\|_p = \lambda^{-n/p}\left\|\sum_{l\in A(\lambda)} f^{(l)}\right\|_p.$$

We note that

$$f^{(l)}(x) = \langle l\rangle^{\frac{n\alpha}{1-\alpha}} \mathrm{e}^{\mathrm{i}x\langle l\rangle^{\frac{\alpha}{1-\alpha}}l} (\mathscr{F}^{-1}\rho)\left(c\langle l\rangle^{\frac{\alpha}{1-\alpha}}(x-C'l)\right).$$

Due to $\mathscr{F}^{-1}\rho$ is a Schwartz function, we see that $|\mathscr{F}^{-1}\rho(x)| \lesssim \langle x\rangle^{-2n}$. We can assume that $\mathscr{F}^{-1}\rho(0)=1$, which follows that there exists a $\delta>0$ such that

$$|\mathscr{F}^{-1}\rho(x)| \geq 1/2, \quad |x|\leq\delta.$$

Hence, for any $k\in A(\lambda)$, $x\in B(C'k, \delta/c\langle k\rangle^{\frac{\alpha}{1-\alpha}})$,

$$\left|\sum_{l\in A(\lambda)} f^{(l)}(x)\right| \geq \frac{\lambda^{-n\alpha}}{2} - C\lambda^{-n\alpha}\sup_{|y|\lesssim\delta}\sum_{l\in\mathbb{Z}^n}\left(y+C'c\,l\right)^{-2n}.$$

We can take $C'c\gg 1$. It follows that for any $k\in A(\lambda)$,

$$\left|\sum_{l\in A(\lambda)} f^{(l)}(x)\right| \geq \frac{\lambda^{-n\alpha}}{2} - C\lambda^{-n\alpha}(C'c)^{-n} \geq \frac{\lambda^{-n\alpha}}{4}, \quad x\in B(C'k, \delta/c\langle k\rangle^{\frac{\alpha}{1-\alpha}}).$$

So, we have

$$\left\|\sum_{l\in A(\lambda)} f^{(l)}(x)\right\|_p \gtrsim \lambda^{-n\alpha}\lambda^{\frac{2n\alpha-n}{p}}.$$

It follows that

$$\|f_\lambda\|^\circ_{M^{s,\alpha}_{p,q}} \gtrsim \lambda^{-n/p+s+n(1-\alpha)(1/q-1/p)}\|f\|_{M^{s,\alpha}_{p,q}}.$$

The result follows. □

Proof of Theorem 4.2. (**Necessity**) **Case 1.** Let us assume that $B^{s_1}_{p,q} \subset M^{s_2}_{p,q}$.

Case 1.1. We show that $s_1 \ge s_2$. Let $k \in \mathbb{Z}^n$, $|k| \gg 1$ with $B(\langle k\rangle^{\frac{\alpha}{1-\alpha}}k, C\langle k\rangle^{\frac{\alpha}{1-\alpha}}) \subset \{\xi : 5 \cdot 2^{j-3} \le |\xi| \le 3 \cdot 2^{j-1}\}$. We see that

$$\left\|\mathscr{F}^{-1}\left(\rho^{\alpha}_k(2\ \cdot)\right)\right\|^{\circ}_{M^{s_2,\alpha}_{p,q}} \geqslant \langle k\rangle^{\frac{s_2}{1-\alpha}}\|\mathscr{F}^{-1}\rho^{\alpha}_k(2\ \cdot)\|_p \gtrsim \langle k\rangle^{\frac{s_2}{1-\alpha}+\frac{n\alpha}{1-\alpha}\left(1-\frac{1}{p}\right)},$$

$$\left\|\mathscr{F}^{-1}\left(\rho^{\alpha}_k(2\ \cdot)\right)\right\|_{B^{s_1}_{p,q}} \leqslant 2^{js_1}\|\mathscr{F}^{-1}\rho^{\alpha}_k(2\ \cdot)\|_p \lesssim \langle k\rangle^{\frac{s_1}{1-\alpha}+\frac{n\alpha}{1-\alpha}\left(1-\frac{1}{p}\right)}.$$

$B^{s_1}_{p,q} \subset M^{s_2,\alpha}_{p,q}$ follows that $s_1 \geqslant s_2$.

Case 1.2. We show that $s_1 \ge s_2 + n(1-\alpha)(1/p + 1/q - 1)$. One has that

$$\|\mathscr{F}^{-1}\varphi_j\|_{B^{s_1}_{p,q}} \lesssim 2^{js_1 + jn\left(1-\frac{1}{p}\right)}. \tag{6.15}$$

Denote

$$A_j = \{k : \ \mathrm{supp}\rho^{\alpha}_k \subset \{\xi : \ 5 \cdot 2^{j-3} \le |\xi| \le 3 \cdot 2^{j-1}\}\}.$$

$$\|\mathscr{F}^{-1}\varphi_j\|^{\circ}_{M^{s_2,\alpha}_{p,q}} \geqslant \left(\sum_{k\in A_j}\langle k\rangle^{\frac{s_2 q}{1-\alpha}}\|\mathscr{F}^{-1}\rho^{\alpha}_k\|^q_p\right)^{\frac{1}{q}} \gtrsim \left(\sum_{k\in A_j}\langle k\rangle^{\frac{s_2 q}{1-\alpha}+\frac{n\alpha}{1-\alpha}\left(1-\frac{1}{p}\right)q}\right)^{\frac{1}{q}}. \tag{6.16}$$

Noticing that $\#A_j \sim O(2^{nj(1-\alpha)})$, we immediately have

$$\|\mathscr{F}^{-1}\varphi_j\|^{\circ}_{M^{s_2,\alpha}_{p,q}} \gtrsim 2^{s_2 j + n\alpha j\left(1-\frac{1}{p}\right) + n(1-\alpha)j\frac{1}{q}}. \tag{6.17}$$

$B^{s_1}_{p,q} \subset M^{s_2,\alpha}_{p,q}$ implies that $s_1 \geqslant s_2 + n(1-\alpha)\left(\frac{1}{p} + \frac{1}{q} - 1\right)$.

Case 1.3. We show that $s_1 \geqslant s_2 + n(1-\alpha)\left(\frac{1}{q} - \frac{1}{p}\right)$. We denote by A_j the set such that for every $k, l \in \mathbb{Z}^n \cap A_j$ $(l \neq \tilde{l})$, $|k - l| \ge C$ and $|k|^{1/(1-\alpha)} \in [5 \cdot 2^{j-3} + C2^{j\alpha}, 3 \cdot 2^{j-1} - C2^{j\alpha}]$. Put

$$f^{(k)} = \tau_k\left(\mathscr{F}^{-1}\rho^{\alpha}_k\right), \qquad f = \sum_{k\in A_j} f^{(k)}. \tag{6.18}$$

Noticing that $\#A_j \sim O(2^{nj(1-\alpha)})$, we have

$$\|f\|^{\circ}_{M^{s_2,\alpha}_{p,q}} = \left(\sum_{k\in A_j}\langle k\rangle^{\frac{s_2 q}{1-\alpha}}\|f^{(k)}\|^q_p\right)^{\frac{1}{q}} \gtrsim 2^{j\left(s_2 + n(1-\alpha)\frac{1}{q} + n\alpha\left(1-\frac{1}{p}\right)\right)}. \tag{6.19}$$

On the other hand,

$$\|f\|_{B^{s_1}_{p,q}} \lesssim 2^{js_1}\left\|\sum_{k\in A_j} f^{(k)}\right\|_p. \tag{6.20}$$

By Plancherel's identity,

$$\|f\|_2 \lesssim \left(\sum_{k\in A_j}\|\rho^{\alpha}_k(2\ \cdot)\|^2_2\right)^{1/2} \lesssim 2^{nj/2}.$$

Moreover, let us observe that

$$f^{(k)} = C\langle k\rangle^{\frac{n\alpha}{1-\alpha}}(\mathscr{F}^{-1}\rho)(C\langle k\rangle^{\frac{n\alpha}{1-\alpha}}(x-k)).$$

Using the same way as in the proof of Case 1.2.3 in Theorem 3.1, we have

$$|f(x)| \lesssim 2^{n\alpha j}.$$

By Hölder's inequality,

$$\|f\|_p \lesssim 2^{jn\alpha(1-2/p)+nj/p}.$$

It follows from $B^{s_1}_{p,q} \subset M^{s_2,\alpha}_{p,q}$ that $s_1 \geqslant s_2 + n(1-\alpha)\big(\frac{1}{q} - \frac{1}{p}\big)$.

Case 2. We assume that $M^{s_1,\alpha_1}_{p,q} \subset B^{s_2}_{p,q}$.

Case 2.1. Let $j \gg 1$. One can find some $k \in \mathbb{Z}^n$ verifying $\rho^\alpha_k \varphi_j = \rho^\alpha_k$. It follows that $2^j \sim \langle k\rangle^{1/(1-\alpha)}$. Thus,

$$\|\mathscr{F}^{-1}\rho^\alpha_k\|_{B^{s_2}_{p,q}} \geqslant 2^{js_2}\|\mathscr{F}^{-1}\rho^\alpha_k\|_p \gtrsim 2^{js_2}\langle k\rangle^{\frac{n\alpha}{1-\alpha}\left(1-\frac{1}{p}\right)}, \tag{6.21}$$

$$\|\mathscr{F}^{-1}\rho^\alpha_k\|^\circ_{M^{s_1,\alpha}_{p,q}} \lesssim \langle k\rangle^{\frac{s_1}{1-\alpha}+\frac{n\alpha}{1-\alpha}\left(1-\frac{1}{p}\right)}. \tag{6.22}$$

So, we have $s_1 \geqslant s_2$.

Case 2.2. Let $j \gg 1$. We have

$$\big\|\mathscr{F}^{-1}\varphi_j\big\|_{B^{s_2}_{p,q}} \geqslant 2^{js_2}\|\mathscr{F}^{-1}\varphi^2_j\|_p \gtrsim 2^{js_2+jn\left(1-\frac{1}{p}\right)}. \tag{6.23}$$

On the other hand,

$$\big\|\mathscr{F}^{-1}\varphi_j\big\|_{M^{s_1,\alpha}_{p,q}} \lesssim \left(\sum_{|k|^{1/(1-\alpha)}\sim 2^j} \langle k\rangle^{\frac{s_1 q}{1-\alpha}} 2^{jn\alpha\left(1-\frac{1}{p}\right)q}\right)^{\frac{1}{q}} \lesssim 2^{jn(1-\alpha)\frac{1}{q}+s_1 j+jn\alpha\left(1-\frac{1}{p}\right)}. \tag{6.24}$$

It follows from $M^{s_1,\alpha}_{p,q} \subset B^{s_2}_{p,q}$ that $s_1 \geqslant s_2 + n(1-\alpha)\big(1 - \frac{1}{p} - \frac{1}{q}\big)$.

Case 2.3. We denote by A_j the set such that for every $k, l \in \mathbb{Z}^n \cap A_j$ $(l \neq k)$, $|k-l| \geq C$ and $|k|^{1/(1-\alpha)} \in [5\cdot 2^{j-3} + C2^{j\alpha},\ 3\cdot 2^{j-1} - C2^{j\alpha}]$. Put

$$f^{(k)} = \tau_{C'k}\mathscr{F}^{-1}\left(\rho\left(\frac{\cdot\, - \langle k\rangle^{\frac{\alpha}{1-\alpha}}k}{c\langle k\rangle^{\frac{\alpha}{1-\alpha}}}\right)\right), \quad f = \sum_{k\in A_j} f^{(k)}. \tag{6.25}$$

Using the same way as in the proof of Case 2.2.3 in Theorem 3.1,

$$\|f\|^\circ_{M^{s_1,\alpha}_{p,q}} \lesssim 2^{j\left[s_1+n(1-\alpha)\frac{1}{q}+n\alpha\left(1-\frac{1}{p}\right)\right]}, \tag{6.26}$$

$$\|f\|_{B^{s_2}_{p,q}} = 2^{js_2}\,\|f\|_p \gtrsim 2^{j\left[s_2+n(1-\alpha)\frac{1}{p}+n\alpha\left(1-\frac{1}{p}\right)\right]}. \tag{6.27}$$

From $M^{s_1,\alpha}_{p,q} \subset B^{s_2}_{p,q}$ it follows that $s_1 \geqslant s_2 + n(1-\alpha)\big(\frac{1}{p} - \frac{1}{q}\big)$. □

Proof of Theorem 4.1. (**Necessity**) We separate the proof into the following two cases.
Case 1. We assume that $M^{s_1,\alpha_1}_{p,q} \subset M^{s_2,\alpha_2}_{p,q}$ with $\alpha_1 \geqslant \alpha_2$.
Case 1.1. We show that $s_1 \geq s_2$. Denote

$$\Lambda_2(k) = \{l \in \mathbb{Z}^n : \ \rho_k^{\alpha_2}\rho_l^{\alpha_1} \neq 0\}.$$

We have

$$\left\|\mathscr{F}^{-1}\rho_k^{\alpha_2}\right\|^\circ_{M^{s_2,\alpha_2}_{p,q}} \geqslant \langle k\rangle^{\frac{s_2}{1-\alpha_2}} \|\mathscr{F}^{-1}\rho_k^{\alpha_2}\rho_k^{\alpha_2}\|_p \gtrsim \langle k\rangle^{\frac{s_2}{1-\alpha_2}+\frac{n\alpha_2}{1-\alpha_2}\left(1-\frac{1}{p}\right)}, \tag{6.28}$$

and

$$\left\|\mathscr{F}^{-1}\rho_k^{\alpha_2}\right\|^\circ_{M^{s_1,\alpha_1}_{p,q}} \lesssim \left(\sum_{l\in\Lambda_2(k)} \langle l\rangle^{\frac{s_1 q}{1-\alpha_1}} \|\mathscr{F}^{-1}\rho_k^{\alpha_2}\rho_l^{\alpha_1}\|_p^q\right)^{\frac{1}{q}}. \tag{6.29}$$

Using Young's inequality and Proposition 2.1, respectively for $p \geq 1$ and $p < 1$, we have

$$\|\mathscr{F}^{-1}\rho_k^{\alpha_2}\rho_l^{\alpha_1}\|_p \lesssim \|\mathscr{F}^{-1}\rho_k^{\alpha_2}\|_p \lesssim \langle k\rangle^{\frac{n\alpha_2}{1-\alpha_2}\left(1-\frac{1}{p}\right)}.$$

Since $\#\Lambda_2(k)$ is finite and $|k|^{\frac{1}{1-\alpha_2}} \sim |l|^{\frac{1}{1-\alpha_1}}$ for all $l \in \lambda_2(k)$, one has that

$$\left\|\mathscr{F}^{-1}\rho_k^{\alpha_2}\right\|^\circ_{M^{s_1,\alpha_1}_{p,q}} \lesssim \left(\sum_{k\in\Lambda_2(k)} \langle k\rangle^{\frac{sq}{1-\alpha_1}} \langle k\rangle^{\frac{n\alpha_2}{1-\alpha_2}\left(1-\frac{1}{p}\right)q}\right)^{\frac{1}{q}} \lesssim \langle k\rangle^{\frac{s}{1-\alpha_2}+\frac{n\alpha_2}{1-\alpha_2}\left(1-\frac{1}{p}\right)}. \tag{6.30}$$

From $M^{s,\alpha_1}_{p,q} \subset M^{s_2,\alpha_2}_{p,q}$ it follows that $s_1 \geqslant s_2$.
Case 1.2. Let $k \in \mathbb{Z}^n$ with $|k| \gg 1$. Denote

$$\Lambda^*(k) = \left\{l \in \mathbb{Z}^n : \rho_k^{\alpha_1}\rho_l^{\alpha_2} = \rho_l^{\alpha_2}\right\}. \tag{6.31}$$

We have

$$\|\mathscr{F}^{-1}\rho_k^{\alpha_1}\|_{M^{s_1,\alpha_1}_{p,q}} \lesssim \langle k\rangle^{\frac{s_1}{1-\alpha_1}+\frac{n\alpha_1}{1-\alpha_1}\left(1-\frac{1}{p}\right)}. \tag{6.32}$$

Since $\#\Lambda^*(k) \sim \langle k\rangle^{\frac{n(\alpha_1-\alpha_2)}{1-\alpha_1}}$ and $\langle k\rangle^{\frac{1}{1-\alpha_1}} \sim \langle l\rangle^{\frac{1}{1-\alpha_2}}$ for all $l \in \Lambda^*(k)$, one has that

$$\|\mathscr{F}^{-1}\rho_k^{\alpha_1}\|^\circ_{M^{s_2,\alpha_2}_{p,q}} \geqslant \left(\sum_{l\in\Lambda^*(k)} \langle l\rangle^{\frac{s_2 q}{1-\alpha_2}} \|\mathscr{F}^{-1}\eta_l^{\alpha_2}\|_p^q\right)^{\frac{1}{q}} \gtrsim \langle k\rangle^{\frac{n(\alpha_1-\alpha_2)}{q(1-\alpha_1)}+\frac{s_2}{1-\alpha_1}+\frac{n\alpha_2}{1-\alpha_1}\left(1-\frac{1}{p}\right)}. \tag{6.33}$$

$M^{s,\alpha_1}_{p,q} \subset M^{s_2,\alpha_2}_{p,q}$ implies that $s_1 \geqslant s_2 + n(\alpha_1-\alpha_2)\left(\frac{1}{p}+\frac{1}{q}-1\right)$.
Case 1.3. We show that $s_1 \geqslant s_2 + n(\alpha_1-\alpha_2)\left(\frac{1}{q}-\frac{1}{p}\right)$. Let $\Lambda_0^*(k)$ be the subset of $\Lambda^*(k)$ such that $|l-\tilde{l}| \geq C$ for all $l, \tilde{l} \in \Lambda_0^*(k)$ $(l \neq \tilde{l})$. Denote

$$f^{(l)} = \tau_l \mathscr{F}^{-1}\rho_l^{\alpha_2}, \qquad f = \sum_{l\in\Lambda_0^*(k)} f^{(l)}. \tag{6.34}$$

It follows that

$$\|f\|^\circ_{M^{s_2,\alpha_2}_{p,q}} = \left(\sum_{l\in\Lambda_0^*(k)} \langle l\rangle^{\frac{s_2 q}{1-\alpha_2}} \|f^{(l)}\|_p^q\right)^{\frac{1}{q}} \gtrsim \langle k\rangle^{\frac{s_2}{1-\alpha_1}+\frac{n(\alpha_1-\alpha_2)}{q(1-\alpha_1)}+\frac{n\alpha_2}{1-\alpha_1}\left(1-\frac{1}{p}\right)}. \tag{6.35}$$

On the other hand,

$$\|f\|_2 = \left\| \sum_{l\in\Lambda_0^*(k)} \rho_l^{\alpha_2} \right\|_2 \sim \langle k\rangle^{\frac{n\alpha_2}{2(1-\alpha_1)}+\frac{n(\alpha_1-\alpha_2)}{2(1-\alpha_1)}},$$

and noticing that $|f^{(l)}(x)| = \langle l\rangle^{\frac{n\alpha_2}{(1-\alpha_2)}}|(\mathscr{F}^{-1})(C\langle l\rangle^{\frac{\alpha_2}{(1-\alpha_2)}}(x-l))|$, we have

$$\|f\|_\infty \lesssim \langle k\rangle^{\frac{n\alpha_2}{(1-\alpha_1)}}.$$

Hence,

$$\|f\|_p \lesssim \langle k\rangle^{\frac{n\alpha_2}{(1-\alpha_1)}(1-\frac{1}{p})+\frac{n(\alpha_1-\alpha_2)}{p(1-\alpha_1)}}.$$

It follows that

$$\|f\|^\circ_{M^{s_1,\alpha_1}_{p,q}} \lesssim \langle k\rangle^{\frac{s_1}{1-\alpha_1}}\|f\|_p \lesssim \langle k\rangle^{\frac{s_1}{1-\alpha_1}+\frac{n(\alpha_1-\alpha_2)}{p(1-\alpha_1)}+\frac{n\alpha_2}{1-\alpha_1}\left(1-\frac{1}{p}\right)}. \tag{6.36}$$

$M^{s,\alpha_1}_{p,q} \subset M^{s_2,\alpha_2}_{p,q}$ implies that $s_1 \geqslant s_2 + n(\alpha_1-\alpha_2)\big(\frac{1}{q}-\frac{1}{p}\big)$.
Case 2. We assume that $M^{s_1,\alpha_1}_{p,q} \subset M^{s_2,\alpha_2}_{p,q}, \alpha_1 < \alpha_2$. The idea is the same as in the proof of Theorem 4.2 and we only give a sketch proof.
Case 2.1. We show that $s_1 \geq s_2$. Let $k \in \mathbb{Z}^n$, $|k| \gg 1$. One can find some $l \in \mathbb{Z}^n$ such that $\rho_k^{\alpha_1}\rho_l^{\alpha_2} = \rho_k^{\alpha_1}$, Then,

$$\|\mathscr{F}^{-1}\rho_k^{\alpha_1}\|^\circ_{M^{s_2,\alpha_2}_{p,q}} \gtrsim \langle k\rangle^{\frac{s_2}{1-\alpha_1}+\frac{n\alpha_1}{1-\alpha_1}\left(1-\frac{1}{p}\right)}, \tag{6.37}$$

$$\|\mathscr{F}^{-1}\rho_k^{\alpha_1}\|^\circ_{M^{s_1,\alpha_1}_{p,q}} \lesssim \langle k\rangle^{\frac{s_1}{1-\alpha_1}+\frac{n\alpha_1}{1-\alpha_1}\left(1-\frac{1}{p}\right)}. \tag{6.38}$$

$M^{s_1,\alpha_1}_{p,q} \subset M^{s_2,\alpha_2}_{p,q}$ implies that $s_1 \geqslant s_2$.
Case 2.2. We show that $s_1 \geqslant s_2 + n(\alpha_1-\alpha_2)\big(\frac{1}{p}+\frac{1}{q}-1\big)$.

$$\big\|\mathscr{F}^{-1}\rho_k^{\alpha_2}\big\|^\circ_{M^{s_2,\alpha_2}_{p,q}} \gtrsim \langle k\rangle^{\frac{s_2}{1-\alpha_2}+\frac{n\alpha_2}{1-\alpha_2}\left(1-\frac{1}{p}\right)}, \tag{6.39}$$

$$\big\|\mathscr{F}^{-1}\rho_k^{\alpha_2}\big\|^\circ_{M^{s_1,\alpha_1}_{p,q}} \lesssim \langle k\rangle^{\frac{n(\alpha_2-\alpha_1)}{q(1-\alpha_2)}+\frac{s_1}{1-\alpha_2}+\frac{n\alpha_1}{1-\alpha_2}\left(1-\frac{1}{p}\right)}. \tag{6.40}$$

$M^{s_1,\alpha_1}_{p,q} \subset M^{s_2,\alpha_2}_{p,q}$ implies that $s_1 \geqslant s_2 + n(\alpha_1-\alpha_2)\big(\frac{1}{p}+\frac{1}{q}-1\big)$.
Case 2.3. Let $k \in \mathbb{Z}^n$ with $|k| \gg 1$, and

$$\Lambda^*(k) = \left\{l \in \mathbb{Z}^n : \rho_l^{\alpha_1}\rho_k^{\alpha_2} = \rho_l^{\alpha_1}\right\}. \tag{6.41}$$

Let $\Lambda_0^*(k)$ be the subset of $\Lambda^*(k)$ such that $|l-\tilde{l}| \geq C$ for all $l, \tilde{l} \in \Lambda_0^*(k)$ $(l \neq \tilde{l})$. It is easy to see that $\#\Lambda_0^*(k) \sim \langle k\rangle^{\frac{n(\alpha_2-\alpha_1)}{1-\alpha_2}}$. Put

$$f^{(l)} = \tau_{Cl}\mathscr{F}^{-1}\left(\rho_l^{\alpha_1}\right), \qquad f = \sum_{l\in\Lambda_0^*(k)} f^{(l)}. \tag{6.42}$$

then,

$$\|f\|^\circ_{M^{s_1,\alpha_1}_{p,q}} \lesssim \langle k\rangle^{\frac{s_1}{1-\alpha_2}+\frac{n(\alpha_2-\alpha_1)}{q(1-\alpha_2)}+\frac{n\alpha_1}{1-\alpha_2}\left(1-\frac{1}{p}\right)}, \tag{6.43}$$

$$\|f\|^\circ_{M^{s_2,\alpha_2}_{p,q}} \gtrsim \langle k\rangle^{\frac{s_2}{1-\alpha_2}+\frac{n(\alpha_2-\alpha_1)}{p(1-\alpha_2)}+\frac{n\alpha_1}{1-\alpha_2}\left(1-\frac{1}{p}\right)}. \tag{6.44}$$

$M^{s_1,\alpha_1}_{p,q} \subset M^{s_2,\alpha_2}_{p,q}$ implies that $s_1 \geqslant s_2 + n(\alpha_2-\alpha_1)\big(\frac{1}{p}-\frac{1}{q}\big)$. $\square$

7 Counterexamples for the algebra structure

In order to show that our results are sharp, we need an α-covering which is a slightly modified version in [9]. Let $Q(a,r) := \prod_{i=1}^{n}[a_i - r, a_i + r]$ and we consider the following covering of $\mathbb{R}$:

$$Q_0 = [-1,1], \quad Q_j = Q(|j|^{\alpha/(1-\alpha)}j,\ r_j|j|^{\alpha/(1-\alpha)}),\ j \neq 0.$$

Lemma 7.1. *Let $r > 1/2(1-\alpha)$. There exists $j_0 \in \mathbb{N}$, such that $\{Q_j\}_{j\in\mathbb{Z}}$ is an α-covering of $\mathbb{R}$, where*

$$r_j = \begin{cases} r, & |j| > j_0, \\ \text{suitable}, & |j| \leq j_0. \end{cases}$$

Moreover, if $r < 8/15(1-\alpha)$, then

$$Q_{j\pm1} \cap Q(|j|^{\alpha/(1-\alpha)}j,\ \frac{7}{8}r_j|j|^{\alpha/(1-\alpha)}) = \varnothing \tag{7.1}$$

for all $j \in \mathbb{Z}$.

Proof. Let $j > 100$. Noticing that $Q_{j+1} \cap Q_j \neq \varnothing$ if and only if

$$|j+1|^{\alpha/(1-\alpha)}(j+1) - r_{j+1}|j+1|^{\alpha/(1-\alpha)} < |j|^{\alpha/(1-\alpha)}j - r_j|j|^{\alpha/(1-\alpha)}. \tag{7.2}$$

In view of mean value theorem, we see that (7.2) is equivalent to

$$\frac{1}{1-\alpha}|j+\theta|^{\alpha/(1-\alpha)} < r_{j+1}|j+1|^{\alpha/(1-\alpha)} + r_j|j|^{\alpha/(1-\alpha)}, \tag{7.3}$$

where $\theta \in (0,1)$. Take $r_{j+1} = r_j = r$. Hence, there exists $j_0 := j_0(\alpha)$ such that for any $j > j_0$, (7.3) holds. Next, if $j > j_0$, we have

$$|j+1|^{\alpha/(1-\alpha)}(j+1) - r|j+1|^{\alpha/(1-\alpha)} > |j|^{\alpha/(1-\alpha)}j + \frac{7}{8}r|j|^{\alpha/(1-\alpha)}, \tag{7.4}$$

which implies (7.1) for $j > j_0$. If $j \leq j_0$, one can choose suitable r_j so that the conclusion holds. □

Using Lemma 7.1 and the idea as in [9], we now construct a new α-covering of $\mathbb{R}^n$, where the original idea goes back to Lizorkin's dyadic decomposition to $\mathbb{R}^n$. Let $j \in \mathbb{Z}$ with $|j| > j_0$. We may assume $8|j|/7r \in \mathbb{N}$. We divide $[-|j|^{\frac{1}{1-\alpha}}, |j|^{\frac{1}{1-\alpha}}]$ into $16|j|/7r$ equal intervals:

$$[-|j|^{\frac{1}{1-\alpha}},\ |j|^{\frac{1}{1-\alpha}}] = [r_{j,-N_j}, r_{j,-N_j+1}] \cup ... \cup [r_{j,N_j-1}, r_{j,N_j}].$$

Denote

$$\mathscr{R} = \{r_{j,s} : j \in \mathbb{N}, s = -N_j, \cdots, N_j\}.$$

We further write

$$\mathscr{K}_j^n = \{k = (k_1, \cdots, k_n) : k_i \in \mathscr{R},\ \max_{1\leq i\leq n}|k_i| = |j|^{\frac{1}{1-\alpha}}\}.$$

For any $k \in \mathscr{K}_j^n$, we write

$$Q_{kj} = Q(k, r|j|^{\frac{\alpha}{1-\alpha}}), \quad |j| > j_0.$$

From the construction of Q_{kj} one sees that

$$\begin{cases} \#\{Q_{k'j'}:\ Q_{k'j'}\cap Q_{kj}\neq\varnothing\}=2n, \\ Q(k,r|j|^{\frac{\alpha}{1-\alpha}}/2)\cap Q_{k'j'}\neq\varnothing \Leftrightarrow k'=k, j'=j. \end{cases} \tag{7.5}$$

For $|j|\le j_0$, one can choose suitable r_j, and in a similar way as above to define

$$Q_{kj}=Q(k,r_j|j|^{\frac{\alpha}{1-\alpha}}),\quad 1\le|j|\le j_0,\quad Q_0=Q(0,r_0)$$

so that

$$\begin{cases} \#\{Q_{k'j'}:\ Q_{k'j'}\cap Q_{kj}\neq\varnothing\}=2n, \\ Q(k,r_j|j|^{\frac{\alpha}{1-\alpha}}/2)\cap Q_{k'j'}\neq\varnothing \Leftrightarrow k'=k, j'=j \quad for\ j\neq 0, \\ Q(0,r_0/2)\cap Q_{k'j'}\neq\varnothing \Leftrightarrow Q_{k'j'}=Q_0. \end{cases} \tag{7.6}$$

Lemma 7.2. *$\{Q_0\}\cup\{Q_{kj}\}_{j\in\mathbb{Z}\backslash\{0\},\ k\in\mathscr{K}_j}$ as in (7.5) and (7.6) is an α-covering of $\mathbb{R}^n$.*

Let $\eta:\mathbb{R}\to[0,1]$ be a smooth bump function satisfying

$$\eta(\xi):=\begin{cases} 1, & |\xi|\le 1/2, \\ \text{smooth}, & 1/2<|\xi|\le 1, \\ 0, & |\xi|\ge 1. \end{cases} \tag{7.7}$$

Let r and r_j be as in (7.5) and (7.6), respectively. Denote for $i=1,\cdots,n$, $j\neq 0$,

$$\phi_{kj}(\xi_i)=\eta\left(\frac{\xi_i-k_i}{r_j|j|^{\frac{\alpha}{1-\alpha}}}\right),\quad k=(k_1,\cdots,k_n)\in\mathscr{K}_j,\quad \phi_0(\xi_i)=\eta\left(\frac{\xi_i}{r_0}\right),$$

$$\phi_{kj}(\xi)=\phi_{kj}(\xi_1)...\phi_{kj}(\xi_n),\quad \phi_0(\xi)=\phi_0(\xi_1)...\phi_0(\xi_n).$$

We put

$$\psi_{kj}(\xi)=\frac{\phi_{kj}(\xi)}{\phi_0(\xi)+\sum_{k\in\mathscr{K}_j,j\in\mathbb{Z}\backslash\{0\}}\phi_{kj}(\xi)},\quad \psi_0(\xi)=\frac{\phi_0(\xi)}{\phi_0(\xi)+\sum_{k\in\mathscr{K}_j,j\in\mathbb{Z}\backslash\{0\}}\phi_{kj}(\xi)}. \tag{7.8}$$

Lemma 7.3. *$\{\psi_0\}\cup\{\psi_{kj}\}_{j\in\mathbb{Z}\backslash\{0\},\ k\in\mathscr{K}_j}$ as in (7.8) is a p-BAPU.*

On the basis of the above *p-BAPU*, we immediately have

Proposition 7.1. *Let $0<\alpha<1$, $0<p,q\leqslant\infty$, then*

$$\|f\|_{M^{s,\alpha}_{p,q}}=\left(\|\mathscr{F}^{-1}\psi_0\mathscr{F}f\|^q_{L^p(\mathbb{R}^n)}+\sum_{j\in\mathbb{Z}\backslash\{0\}}\langle j\rangle^{sq/(1-\alpha)}\sum_{k\in\mathscr{K}_j}\|\mathscr{F}^{-1}\psi_{kj}\mathscr{F}f\|^q_{L^p(\mathbb{R}^n)}\right)^{1/q}$$

is an equivalent norm on α-modulation space.

Theorem 7.1. *Let $0\le\alpha<1$, $(1/p,1/q)\in D_2$, $p>1$. If $s<s_0$, then $M^{s,\alpha}_{p,q}$ is not a Banach algebra.*

Proof. **Step 1. $p > 1$, $q \le 1$.** Let χ_A be the characteristic function on A. Now we take for $J \gg j_0$, $\ell = (J, J, ..., J)$,

$$\widehat{f} = \chi_{A(J)}, \quad \widehat{g} = \chi_{-A(J)}, \quad A(J) = Q(|J|^{\alpha/(1-\alpha)}\ell,\ r|J|^{\alpha/(1-\alpha)}/2). \tag{7.9}$$

Noticing that

$$(\chi_{[B-b,B+b]} * \chi_{[-B-b,-B+b]})(\xi) = \begin{cases} 0, & |\xi| \ge 2b, \\ 2b - |\xi|, & |\xi| < 2b. \end{cases} \tag{7.10}$$

Hence,

$$\left(\chi_{A(J)} * \chi_{-A(J)}\right)(\xi) = \begin{cases} 0, & |\xi_i| \ge r|J|^{\frac{\alpha}{(1-\alpha)}} \text{ for some } i = 1, ..., n, \\ \prod_{i=1}^{n}(r|J|^{\frac{\alpha}{(1-\alpha)}} - |\xi_i|), & |\xi_i| < r|J|^{\frac{\alpha}{(1-\alpha)}} \text{ for all } i = 1, ..., n. \end{cases} \tag{7.11}$$

Hence,

$$\mathrm{supp}\widehat{f} * \widehat{g} = \{\xi \in \mathbb{R}^n : |\xi_i| \le r|J|^{\frac{\alpha}{(1-\alpha)}},\ i = 1, ..., n\}.$$

One has that

$$\begin{aligned} \prod_{i=1}^{n}(r|J|^{\frac{\alpha}{(1-\alpha)}} + \xi_i) &= (r|J|^{\frac{\alpha}{(1-\alpha)}})^n + (r|J|^{\frac{\alpha}{(1-\alpha)}})^{n-1}\sum_i \xi_i \\ &\quad + (r|J|^{\frac{\alpha}{(1-\alpha)}})^{n-2}\sum_{i<j}\xi_i\xi_j + ... + \xi_1...\xi_n \\ &:= (r|J|^{\frac{\alpha}{(1-\alpha)}})^n + R(\xi, J). \end{aligned}$$

Let us write

$$\mathscr{A}_j = \{k \in \mathscr{K}_j^n : \ \text{ if } \ \xi, \eta \in \mathrm{supp}\psi_{k,j}, \text{ then } \xi_i\eta_i > 0 \ \text{ for all } \ i = 1, ..., n\}.$$

Let $1 \ll j < \varepsilon|J|^\alpha$ $(0 < \varepsilon \ll 1)$ and $k \in \mathscr{A}_j$. We may assume that $\xi_i > 0$ if $\xi \in \mathrm{supp}\psi_{kj}$. Noticing that $\mathrm{supp}\psi_{kj} \subset Q(0, r|J|^{\alpha/(1-\alpha)})$, we have

$$\begin{aligned} \|\mathscr{F}^{-1}\psi_{kj}\mathscr{F}(fg)\|_p &= \left\|\mathscr{F}^{-1}\psi_{kj}\prod_{i=1}^{n}(r|J|^{\frac{\alpha}{(1-\alpha)}} - \xi_i)\right\|_p \\ &\ge (r|J|^{\frac{\alpha}{1-\alpha}})^n\|\mathscr{F}^{-1}\psi_{kj}\|_p - \left\|\mathscr{F}^{-1}(\psi_{kj}R(\xi, J))\right\|_p \\ &\ge c|J|^{\frac{n\alpha}{1-\alpha}}|j|^{\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})} - \left\|\mathscr{F}^{-1}(\psi_{kj}R(\xi, J))\right\|_p. \end{aligned} \tag{7.12}$$

$$\begin{aligned} \left\|\mathscr{F}^{-1}(\psi_{kj}R(\xi, J))\right\|_p &\lesssim |J|^{\frac{(n-1)\alpha}{1-\alpha}}\left\|\mathscr{F}^{-1}(\psi_{kj}\sum_{i=1}^{n}\xi_i)\right\|_p \\ &\quad + |J|^{\frac{(n-2)\alpha}{1-\alpha}}\left\|\mathscr{F}^{-1}(\psi_{kj}\sum_{i<j}\xi_i\xi_j)\right\|_p + ... + \left\|\mathscr{F}^{-1}(\psi_{kj}\xi_1...\xi_n)\right\|_p. \end{aligned} \tag{7.13}$$

For instance, we estimate $\|\mathscr{F}^{-1}(\psi_{kj}\xi_1\xi_2)\|_p$. Let k be the center of suppψ_{kj}. We have

$$\begin{aligned}\left\|\mathscr{F}^{-1}(\psi_{kj}\xi_1\xi_2)\right\|_p &\lesssim |k_1||k_2| \left\|\mathscr{F}^{-1}\psi_{kj}\right\|_p + \left\|\mathscr{F}^{-1}(\psi_{kj}(\xi_1-k_1)(\xi_2-k_2))\right\|_p \\ &\quad + |k_1| \left\|\mathscr{F}^{-1}(\psi_{kj}(\xi_2-k_2))\right\|_p + |k_2| \left\|\mathscr{F}^{-1}(\psi_{kj}(\xi_1-k_1))\right\|_p \\ &\lesssim |j|^{\frac{2}{1-\alpha}}|j|^{\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})} \lesssim \varepsilon^2|J|^{\frac{2\alpha}{1-\alpha}}|j|^{\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})}. \end{aligned} \tag{7.14}$$

So, one has that

$$\left\|\mathscr{F}^{-1}(\psi_{kj}R(\xi,J))\right\|_p \lesssim \varepsilon|J|^{\frac{n\alpha}{1-\alpha}}|j|^{\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})}. \tag{7.15}$$

It follows from (7.12) and (7.15) that

$$\|\mathscr{F}^{-1}\psi_{kj}\mathscr{F}(fg)\|_p \gtrsim |J|^{\frac{n\alpha}{1-\alpha}}|j|^{\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})}. \tag{7.16}$$

(7.23) yields

$$\begin{aligned}\|fg\|_{M^{s,\alpha}_{p,q}} &\gtrsim \left(\sum_{1\ll j\le\varepsilon|J|^\alpha}|j|^{\frac{sq}{1-\alpha}}\sum_{k\in\mathscr{A}_j}\|\mathscr{F}^{-1}\psi_{kj}\mathscr{F}(fg)\|_p^q\right)^{1/q} \\ &\gtrsim |J|^{\frac{n\alpha}{1-\alpha}}\left(\sum_{1\ll j\le\varepsilon|J|^\alpha}|j|^{\frac{sq}{1-\alpha}}\sum_{k\in\mathscr{A}_j}|j|^{\frac{n\alpha q}{1-\alpha}(1-\frac{1}{p})}\right)^{1/q} \\ &\gtrsim |J|^{\frac{n\alpha}{(1-\alpha)}+\alpha\left(\frac{s}{(1-\alpha)}+\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})+\frac{n}{q}\right)}. \end{aligned} \tag{7.17}$$

On the other hand,

$$\|f\|_{M^{s,\alpha}_{p,q}} \sim |J|^{\frac{s}{(1-\alpha)}}\|\mathscr{F}^{-1}\chi_{A(J)}\|_p \sim |J|^{\frac{s}{(1-\alpha)}+\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})}. \tag{7.18}$$

Similarly,

$$\|g\|_{M^{s,\alpha}_{p,q}} \sim |J|^{\frac{s}{(1-\alpha)}+\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})}. \tag{7.19}$$

Hence, in order to $M^{s,\alpha}_{p,q}$ forms an algebra, one must has that

$$\frac{2s}{(1-\alpha)}+\frac{2n\alpha}{1-\alpha}(1-\frac{1}{p}) \ge \frac{n\alpha}{(1-\alpha)}+\alpha\left(\frac{s}{(1-\alpha)}+\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})+\frac{n}{q}\right). \tag{7.20}$$

Namely,

$$s \ge \frac{n\alpha}{p}+\frac{n\alpha(1-\alpha)}{(2-\alpha)}\left(\frac{1}{q}-1\right). \tag{7.21}$$

Step 2. $(1/p,1/q)\in[0,1]^2\cap D_2$. Let $J\gg 1$. Put

$$\widehat{f}(\xi)=\chi_{[J^{1/(1-\alpha)},3J^{1/(1-\alpha)}]^n}(\xi), \quad \widehat{g}(\xi)=\chi_{[-3J^{1/(1-\alpha)},-J^{1/(1-\alpha)}]^n}(\xi).$$

In view of (7.10) we have

$$(\widehat{f}*\widehat{g})(\xi)=\begin{cases} 0, & |\xi_i|\ge 2J^{\frac{1}{(1-\alpha)}} \text{ for some } i=1,...,n, \\ \prod_{i=1}^n(2J^{\frac{1}{(1-\alpha)}}-|\xi_i|), & |\xi_i|\le 2J^{\frac{1}{(1-\alpha)}} \text{ for all } i=1,...,n. \end{cases} \tag{7.22}$$

Hence,

$$\operatorname{supp}(\widehat{f} * \widehat{g}) \subset \{\xi : \ |\xi_i| \leq 2J^{\frac{1}{(1-\alpha)}} \quad i = 1, ..., n\}.$$

Using the same way as in (7.23), we have for $|j| \leq \varepsilon J$ and $k \in \mathscr{A}_j$,

$$\|\mathscr{F}^{-1}\psi_{kj}\mathscr{F}(fg)\|_p \gtrsim J^{\frac{n}{1-\alpha}}|j|^{\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})}. \tag{7.23}$$

(7.23) implies that

$$\begin{aligned}\|fg\|_{M^{s,\alpha}_{p,q}} &\gtrsim \left(\sum_{1\ll j\leq \varepsilon J} |j|^{\frac{sq}{1-\alpha}} \sum_{k\in\mathscr{A}_j} \|\mathscr{F}^{-1}\psi_{kj}\mathscr{F}(fg)\|_p^q\right)^{1/q} \\ &\gtrsim J^{\frac{n}{1-\alpha}} \left(\sum_{1\ll j\leq \varepsilon J} |j|^{\frac{sq}{1-\alpha}} \sum_{k\in\mathscr{A}_j} |j|^{\frac{n\alpha q}{1-\alpha}(1-\frac{1}{p})}\right)^{1/q} \\ &\gtrsim J^{\frac{n}{1-\alpha}+\frac{s}{1-\alpha}+\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})+\frac{n}{q}}. \end{aligned} \tag{7.24}$$

On the other hand,

$$\begin{aligned}\|f\|_{M^{s,\alpha}_{p,q}} &\lesssim \left(\sum_{|j|\sim J} |j|^{\frac{sq}{1-\alpha}} \sum_{k\in\mathscr{K}_j} \|\mathscr{F}^{-1}\psi_{kj}\widehat{f}\|_p^q\right)^{1/q} \\ &\lesssim J^{\frac{s}{1-\alpha}+\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})+\frac{n}{q}}. \end{aligned} \tag{7.25}$$

Similarly,

$$\|g\|_{M^{s,\alpha}_{p,q}} \lesssim J^{\frac{s}{1-\alpha}+\frac{n\alpha}{1-\alpha}(1-\frac{1}{p})+\frac{n}{q}}. \tag{7.26}$$

Hence, in order to $M^{s,\alpha}_{p,q}$ forms an algebra, one must has that

$$s \geq \frac{n\alpha}{p} + n(1-\alpha)\left(1-\frac{1}{q}\right). \tag{7.27}$$

□

Acknowledgement. The first named author is supported in part by the National Science Foundation of China, grant 11026053. Part of the work was carried out while the second named author was visiting the Beijing International Center for Mathematical Research (BICMR), he would like to thank BICMR for its hospitality.

References

[1] L. Borup, M. Nielsen, Banach frames for multivariate α-modulation spaces, J. Math. Anal. Appl. **321** (2006), no. 2, 880–895.

[2] L. Borup, M. Nielsen, Boundedness for pseudodifferential operators on multivariate *alpha*-modulation spaces, Ark. Mat. **44** (2006) 241–259.

[3] L. Borup, M. Nielsen, Nonlinear approximation in α-modulation spaces. Math. Nachr. **279** (2006), no. 1-2, 101–120.

[4] A. P. Calderón, Intermediate spaces and interpolation, the complex method, Studia Math., **24** (1964), 113–190.

[5] A. P. Calderón and A. Torchinsky, Parabolic maximal functions associated with a distribution, I, Advances in Math., **16** (1975), 1–64.

[6] A. P. Calderón and A. Torchinsky, Parabolic maximal functions associated with a distribution, II, Advances in Math., **24** (1977), 101–171.

[7] S. Dahlke, M. Fornasier, H. Rauhut, G. Steidl, G. Teschke, Generalized coorbit theory, Banach frames, and the relation to α-modulation spaces, Proc. Lond. Math. Soc., **96** (2008), no. 2, 464–506.

[8] H. G. Feichtinger, *Modulation spaces on locally compact Abelian group*, Technical Report, University of Vienna, 1983.

[9] H. G. Feichtinger, C. Y. Huang, B. X. Wang, Trace operators for modulation, α-modulation and Besov spaces, Appl. Comput. Harmon. Anal., **30** (2011), 110–127.

[10] M. Fornasier, Banach frames for α-modulation spaces, Appl. Comput. Harmon. Anal. **22** (2007), no. 2, 157–175.

[11] P. Gröbner, *Banachräume Glatter Funktionen and Zerlegungsmethoden*, Doctoral thesis, University of Vienna, 1992.

[12] K. Gröchenig, *Foundations of Time-Frequency Analysis*, Birkhäuser Boston, MA, 2001.

[13] M. Kobayashi, M. Sugimoto, N. Tomita, On the L^2-boundedness of pseudo-differential operators and their commutators with symbols in α-modulation spaces. J. Math. Anal. Appl. **350** (2009), no. 1, 157–169.

[14] M. Kobayashi, M. Sugimoto, N. Tomita, Trace ideals for pseudo-differential operators and their commutators with symbols in α-modulation spaces. J. Anal. Math. **107** (2009), 141–160.

[15] M. Sugimoto, N. Tomita, *The dilation property of modulation space and their inclusion relation with Besov spaces*, J. Funct. Anal., **248** (2007), 79–106.

[16] J. Toft, *Continuity properties for modulation spaces, with applications to pseudo-differential calculus, I*, J. Funct. Anal., **207** (2004), 399–429.

[17] J. Toft and P. Wahlberg, Embeddings of α-modulation spaces, Arxiv: 1110.2681.

[18] H. Triebel, *Theory of Function Spaces*, Birkhäuser-Verlag, Basel, 1983.

[19] B. X. Wang and H. Hudzik, The global Cauchy problem for the NLS and NLKG with small rough data, J. Differential Equations, **232** (2007), 36–73.

[20] B. X. Wang and C. Y. Huang, Frequency-uniform decomposition method for the generalized BO, KdV and NLS equations, J. Differential Equations, **239** (2007), 213–250.

[21] B. X. Wang L. Zhao, B. Guo, Isometric decomposition operators, function spaces $E^{\lambda}_{p,q}$ and applications to nonlinear evolution equations. J. Funct. Anal. **233** (2006), no. 1, 1–39.